

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 August 2002 (01.08.2002)

PCT

(10) International Publication Number  
**WO 02/058685 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 31/00**

(21) International Application Number: PCT/US02/02004

(22) International Filing Date: 25 January 2002 (25.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/264,275 26 January 2001 (26.01.2001) US  
60/323,842 21 September 2001 (21.09.2001) US

(71) Applicant: **SCHERING CORPORATION** [US/US];  
Patent Department - K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: **DAVIS, Harry, R.**; 64 River Bend Road,  
Berkeley Heights, NJ 07922 (US). **KOSOGLOU, Teddy**;  
2457 Primrose Court, Jamison, Bucks County, PA  
18929-1178 (US).

(74) Agent: **CANNONI, Ann, Marie**; Schering-Plough Cor-  
poration, Patent Department K-6-1 1990, 2000 Galloping  
Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,  
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,  
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA,  
UZ, VN, YU, ZA, ZM.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— *as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations*

**Published:**

— *without international search report and to be republished  
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*



**WO 02/058685 A2**

(54) Title: COMBINATIONS OF NICOTINIC ACID AND DERIVATIVES THEREOF AND STEROL ABSORPTION INHIBITOR(S) AND TREATMENTS FOR VASCULAR INDICATIONS

(57) Abstract: The present invention provides compositions, therapeutic combinations and methods including: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one substituted azetidinone or substituted b-lactam sterol absorption inhibitor which can be useful for treating vascular conditions, diabetes, obesity and lowering plasma levels of sterols.

5                   **COMBINATIONS OF NICOTINIC ACID AND DERIVATIVES THEREOF AND  
STEROL ABSORPTION INHIBITOR(S) AND TREATMENTS FOR VASCULAR  
INDICATIONS**

**CROSS-REFERENCE TO RELATED APPLICATIONS**

                  This application claims the benefit of priority from U.S. Provisional Patent  
Application Serial No. 60/264,275 filed January 26, 2001 and U.S. Provisional Patent  
10   Application Serial No. 60/323,842 filed September 21, 2001, each of which is  
incorporated herein by reference.

**FIELD OF THE INVENTION**

                  The present invention relates to compositions and therapeutic combinations  
15   comprising nicotinic acid or derivatives thereof and certain sterol absorption inhibitors  
for treating hyperlipidemic conditions such as are associated with atherosclerosis,  
hypercholesterolemia and other vascular conditions in mammals.

**BACKGROUND OF THE INVENTION**

20               Atherosclerotic coronary heart disease (CHD) represents the major cause for  
death and vascular morbidity in the western world. Risk factors for atherosclerotic  
coronary heart disease include hypertension, diabetes mellitus, family history, male  
gender, cigarette smoke and serum cholesterol. A total cholesterol level in excess of  
225-250 mg/dl is associated with significant elevation of risk of CHD.

25               Cholesteryl esters are a major component of atherosclerotic lesions and the  
major storage form of cholesterol in arterial wall cells. Formation of cholesteryl esters  
is also a key step in the intestinal absorption of dietary cholesterol. Thus, inhibition of  
cholesteryl ester formation and reduction of serum cholesterol can inhibit the  
progression of atherosclerotic lesion formation, decrease the accumulation of  
30   cholesteryl esters in the arterial wall, and block the intestinal absorption of dietary  
cholesterol.

                  The regulation of whole-body cholesterol homeostasis in mammals and  
animals involves the regulation of dietary cholesterol and modulation of cholesterol  
biosynthesis, bile acid biosynthesis and the catabolism of the cholesterol-containing  
35   plasma lipoproteins. The liver is the major organ responsible for cholesterol  
biosynthesis and catabolism and, for this reason, it is a prime determinant of plasma

cholesterol levels. The liver is the site of synthesis and secretion of very low density lipoproteins (VLDL) which are subsequently metabolized to low density lipoproteins (LDL) in the circulation. LDL are the predominant cholesterol-carrying lipoproteins in the plasma and an increase in their concentration is correlated with increased  
5 atherosclerosis. When intestinal cholesterol absorption is reduced, by whatever means, less cholesterol is delivered to the liver. The consequence of this action is decreased hepatic lipoprotein (VLDL) production and an increase in the hepatic clearance of plasma cholesterol, mostly as LDL. Thus, the net effect of inhibiting intestinal cholesterol absorption is a decrease in plasma cholesterol levels.

10 Nicotinic acid and its derivatives can inhibit hepatic production of VLDL and its metabolite LDL and increases HDL and apo A-1 levels.

U.S. Patents Nos. 5,767,115, 5,624,920, 5,668,990, 5,656,624 and 5,688,787, respectively, disclose hydroxy-substituted azetidinone compounds and substituted  $\beta$ -lactam compounds useful for lowering cholesterol and/or in inhibiting the formation of  
15 cholesterol-containing lesions in mammalian arterial walls. U.S. Patents Nos. 5,846,966 and 5,661,145, respectively, disclose hydroxy-substituted azetidinone compounds or substituted  $\beta$ -lactam compounds in combination with HMG CoA reductase inhibitors for preventing or treating atherosclerosis and reducing plasma cholesterol levels.

20 PCT Patent Application No. WO 00/38725 discloses cardiovascular therapeutic combinations including an ileal bile acid transport inhibitor or cholesteryl ester transport protein inhibitor in combination with a fibric acid derivative, nicotinic acid derivative, microsomal triglyceride transfer protein inhibitor, cholesterol absorption antagonist, phytosterol, stanol, antihypertensive agent or bile acid sequestrant.


25 U.S. Patent No. 5,698,527 discloses ergostanone derivatives substituted with disaccharides as cholesterol absorption inhibitors, employed alone or in combination with certain other cholesterol lowering agents, which are useful in the treatment of hypercholesterolemia and related disorders.

Despite recent improvements in the treatment of vascular disease, there  
30 remains a need in the art for improved compositions and treatments for hyperlipidaemia, atherosclerosis and other vascular conditions.

#### SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising:

(a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (I):



(I)

5 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (I) or of the isomers thereof, or prodrugs of the compounds of Formula (I) or of the isomers, salts or solvates thereof, wherein in Formula (I) above:

Ar¹ and Ar² are independently selected from the group consisting of aryl and R⁴-substituted aryl;

Ar³ is aryl or R⁵-substituted aryl;

X, Y and Z are independently selected from the group consisting of -CH₂-, -CH(lower alkyl)- and -C(dilower alkyl)-;

R and R² are independently selected from the group consisting of -OR⁶, -O(CO)R⁶, -O(CO)OR⁹ and -O(CO)NR⁶R⁷;

R¹ and R³ are independently selected from the group consisting of hydrogen, lower alkyl and aryl;

q is 0 or 1;

r is 0 or 1;

20 m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;


R⁴ is 1-5 substituents independently selected from the group consisting of lower alkyl, -OR⁶, -O(CO)R⁶, -O(CO)OR⁹, -O(CH₂)₁-₅OR⁶, -O(CO)NR⁶R⁷, -NR⁶R⁷, -NR⁶(CO)R⁷, -NR⁶(CO)OR⁹, -NR⁶(CO)NR⁷R⁸, -NR⁶SO₂R⁹, -COOR⁶, -CONR⁶R⁷, -COR⁶, -SO₂NR⁶R⁷, S(O)₀-₂R⁹, -O(CH₂)₁-₁₀-COOR⁶, -O(CH₂)₁-₁₀CONR⁶R⁷, -(lower alkylene)COOR⁶, -CH=CH-COOR⁶, -CF₃, -CN, -NO₂ and halogen;

- $R^5$  is 1-5 substituents independently selected from the group consisting of -  
 $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
 $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ , -  
 $COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 5  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of  
 hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.


- In another embodiment, there is provided a composition comprising: (a) at least  
 10 one of nicotinic acid or derivatives thereof; and (b) a compound represented by  
 Formula (II) below:



(II)

- or pharmaceutically acceptable salt or solvate thereof, or prodrug of the compound of  
 15 Formula (II) or of the salt or solvate thereof.

In another embodiment, the present invention provides a composition  
 comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one  
 sterol absorption inhibitor represented by Formula (III):



(III)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (III) or of the isomers thereof, or prodrugs of the compounds of Formula (III) or of the isomers, salts or solvates thereof, wherein, in Formula (III) above:

$Ar^1$  is  $R^3$ -substituted aryl;  $Ar^2$  is  $R^4$ -substituted aryl;  $Ar^3$  is  $R^5$ -substituted aryl;

5 Y and Z are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(\text{lower alkyl})-$  and  $-C(\text{dilower alkyl})-$ ;

A is selected from  $-O-$ ,  $-S-$ ,  $-S(O)-$  or  $-S(O)_2-$ ;

$R^1$  is selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$  and  $-O(CO)NR^6R^7$ ;  $R^2$  is selected from the group consisting of hydrogen, lower alkyl and  
10 aryl; or  $R^1$  and  $R^2$  together are  $=O$ ;

q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;


$R^5$  is 1-3 substituents independently selected from the group consisting of -  
 $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^9$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
 15  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2\text{-lower alkyl}$ ,  
 $-NR^6SO_2\text{-aryl}$ ,  $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}\text{-alkyl}$ ,  $S(O)_{0-2}\text{-aryl}$ ,  
 $-O(CH_2)_{1-10}\text{-COOR}^6$ ,  $-O(CH_2)_{1-10}\text{CONR}^6R^7$ , o-halogeno, m-halogeno,  
 o-lower alkyl, m-lower alkyl,  $-(\text{lower alkylene})\text{-COOR}^6$ , and  $-\text{CH}=\text{CH}\text{-COOR}^6$ ;

$R^3$  and  $R^4$  are independently 1-3 substituents independently selected from the  
 20 group consisting of  $R^5$ , hydrogen, p-lower alkyl, aryl,  $-\text{NO}_2$ ,  $-\text{CF}_3$  and p-halogeno;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.

25 In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (IV):



(IV)


or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 5 of Formula (IV) or of the isomers thereof, or prodrugs of the compounds of Formula  
 (IV) or of the isomers, salts or solvates thereof, wherein, in Formula (IV) above:

A is selected from the group consisting of R<sup>2</sup>-substituted heterocycloalkyl, R<sup>2</sup>-  
 substituted heteroaryl, R<sup>2</sup>-substituted benzofused heterocycloalkyl, and R<sup>2</sup>-substituted  
 benzofused heteroaryl;

10 Ar<sup>1</sup> is aryl or R<sup>3</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the



R<sup>1</sup> is selected from the group consisting of:


15 -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can  
 also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-G-(CH<sub>2</sub>)<sub>r</sub>-, wherein G is -O-, -C(O)-, phenylene, -NR<sup>8</sup>- or  
 -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-; and

20 -(CH<sub>2</sub>)<sub>f</sub>V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5,  
 provided that the sum of f and g is 1-6;

R<sup>5</sup> is selected from:




R<sup>6</sup> and R<sup>7</sup> are independently selected from the group consisting of  
 25 -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl), -CH=CH- and

$-C(C_1-C_6 \text{ alkyl})=CH-$ ; or  $R^5$  together with an adjacent  $R^6$ , or  $R^5$  together with an adjacent  $R^7$ , form a  $-CH=CH-$  or a  $-CH=C(C_1-C_6 \text{ alkyl})-$  group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when  $R^6$  is  $-CH=CH-$  or  $-C(C_1-C_6 \text{ alkyl})=CH-$ , a is 1; provided that when  $R^7$  is  $-CH=CH-$  or  $-C(C_1-C_6 \text{ alkyl})=CH-$ , b is 1; provided that when a is 2 or 3, the  $R^6$ 's can be the same or different; and provided that when b is 2 or 3, the  $R^7$ 's can be the same or different;

and when Q is a bond,  $R^1$  also can be selected from:



where M is  $-O-$ ,  $-S-$ ,  $-S(O)-$  or  $-S(O)_2-$ ;

X, Y and Z are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(C_1-C_6 \text{ alkyl})-$  and  $-C(\text{di-}(C_1-C_6 \text{ alkyl}))-$ ;

$R^{10}$  and  $R^{12}$  are independently selected from the group consisting of  $-OR^{14}$ ,  $-O(CO)R^{14}$ ,  $-O(CO)OR^{16}$  and  $-O(CO)NR^{14}R^{15}$ ;

$R^{11}$  and  $R^{13}$  are independently selected from the group consisting of hydrogen,  $(C_1-C_6)\text{alkyl}$  and aryl; or  $R^{10}$  and  $R^{11}$  together are  $=O$ , or  $R^{12}$  and  $R^{13}$  together are  $=O$ ;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;


v is 0 or 1;


j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

$R^2$  is 1-3 substituents on the ring carbon atoms selected from the group consisting of hydrogen,  $(C_1-C_{10})\text{alkyl}$ ,  $(C_2-C_{10})\text{alkenyl}$ ,  $(C_2-C_{10})\text{alkynyl}$ ,  $(C_3-C_6)\text{cycloalkyl}$ ,  $(C_3-C_6)\text{cycloalkenyl}$ ,  $R^{17}$ -substituted aryl,  $R^{17}$ -substituted benzyl,  $R^{17}$ -substituted benzyloxy,  $R^{17}$ -substituted aryloxy, halogeno,  $-NR^{14}R^{15}$ ,  $NR^{14}R^{15}(C_1-C_6 \text{ alkylene})-$ ,  $NR^{14}R^{15}C(O)(C_1-C_6 \text{ alkylene})-$ ,  $-NHC(O)R^{16}$ ,  $OH$ ,



$C_1-C_6$  alkoxy,  $-OC(O)R^{16}$ ,  $-COR^{14}$ , hydroxy( $C_1-C_6$ )alkyl,  
 $(C_1-C_6)$ alkoxy( $C_1-C_6$ )alkyl,  $NO_2$ ,  $-S(O)_{0.2}R^{16}$ ,  $-SO_2NR^{14}R^{15}$  and  
 $-(C_1-C_6 \text{ alkylene})COOR^{14}$ ; when  $R^2$  is a substituent on a heterocycloalkyl ring,  $R^2$  is

as defined, or is  $=O$  or ; and, where  $R^2$  is a substituent on a  
5 substitutable ring nitrogen, it is hydrogen, ( $C_1-C_6$ )alkyl, aryl, ( $C_1-C_6$ )alkoxy, aryloxy,  
 $(C_1-C_6)$ alkylcarbonyl, arylcarbonyl, hydroxy,  $-(CH_2)_{1-6}CONR^{18}R^{18}$ ,



wherein J is  $-O-$ ,  $-NH-$ ,  $-NR^{18}-$  or  $-CH_2-$ ;

$R^3$  and  $R^4$  are independently selected from the group consisting of 1-3  
10 substituents independently selected from the group consisting of ( $C_1-C_6$ )alkyl,  
 $-OR^{14}$ ,  $-O(CO)R^{14}$ ,  $-O(CO)OR^{16}$ ,  $-O(CH_2)_{1-5}OR^{14}$ ,  $-O(CO)NR^{14}R^{15}$ ,  $-NR^{14}R^{15}$ ,  
 $-NR^{14}(CO)R^{15}$ ,  $-NR^{14}(CO)OR^{16}$ ,  $-NR^{14}(CO)NR^{15}R^{19}$ ,  $-NR^{14}SO_2R^{16}$ ,  $-COOR^{14}$ ,  
 $-CONR^{14}R^{15}$ ,  $-COR^{14}$ ,  $-SO_2NR^{14}R^{15}$ ,  $S(O)_{0.2}R^{16}$ ,  $-O(CH_2)_{1-10}-COOR^{14}$ ,  
 $-O(CH_2)_{1-10}CONR^{14}R^{15}$ ,  $-(C_1-C_6 \text{ alkylene})-COOR^{14}$ ,  $-CH=CH-COOR^{14}$ ,  $-CF_3$ ,  $-CN$ ,  $-$   
15  $NO_2$  and halogen;

$R^8$  is hydrogen, ( $C_1-C_6$ )alkyl, aryl ( $C_1-C_6$ )alkyl,  $-C(O)R^{14}$  or  $-COOR^{14}$ ;

$R^9$  and  $R^{17}$  are independently 1-3 groups independently selected from the  
group consisting of hydrogen, ( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxy,  $-COOH$ ,  $NO_2$ ,  
 $-NR^{14}R^{15}$ ,  $OH$  and halogeno;


20  $R^{14}$  and  $R^{15}$  are independently selected from the group consisting of hydrogen,  
 $(C_1-C_6)$ alkyl, aryl and aryl-substituted ( $C_1-C_6$ )alkyl;

$R^{16}$  is ( $C_1-C_6$ )alkyl, aryl or  $R^{17}$ -substituted aryl;

$R^{18}$  is hydrogen or ( $C_1-C_6$ )alkyl; and

$R^{19}$  is hydrogen, hydroxy or ( $C_1-C_6$ )alkoxy.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (V):



(V)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (V) or of the isomers thereof, or prodrugs of the compounds of Formula (V) or of the isomers, salts or solvates thereof, wherein, in Formula (V) above:

- 10  $Ar^1$  is aryl,  $R^{10}$ -substituted aryl or heteroaryl;  
 $Ar^2$  is aryl or  $R^4$ -substituted aryl;  
 $Ar^3$  is aryl or  $R^5$ -substituted aryl;  
X and Y are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(\text{lower alkyl})-$  and  $-C(\text{dilower alkyl})-$ ;
- 15 R is  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$  or  $-O(CO)NR^6R^7$ ;  $R^1$  is hydrogen, lower alkyl or aryl; or R and  $R^1$  together are  $=O$ ;  
q is 0 or 1;  
r is 0, 1 or 2;  
m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and
- 20 q is 1, 2, 3, 4 or 5;  
 $R^4$  is 1-5 substituents independently selected from the group consisting of lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}COOR^6$ ,  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;
- 25  $R^5$  is 1-5 substituents independently selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,


-NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, halogen, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;

R<sup>9</sup> is lower alkyl, aryl or aryl-substituted lower alkyl; and

R<sup>10</sup> is 1-5 substituents independently selected from the group consisting of lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (VI):



(VI)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VI) or of the isomers thereof, or prodrugs of the compounds of Formula (VI) or of the isomers, salts or solvates thereof, wherein:

R<sub>1</sub> is

$\overset{|}{\text{CH-}}$ ,  $\overset{|}{\text{C}}(\text{lower alkyl})\text{-}$ ,  $\overset{|}{\text{CF-}}$ ,  $\overset{|}{\text{C}}(\text{OH})\text{-}$ ,  $\overset{|}{\text{C}}(\text{C}_6\text{H}_5)\text{-}$ ,  $\overset{|}{\text{C}}(\text{C}_6\text{H}_4\text{-R}_{15})\text{-}$ ,  
 $\overset{|}{\text{N-}}$  or  $\overset{|}{\text{N}}\text{O}^-$ ;

R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of:

-CH<sub>2</sub>-, -CH(lower alkyl)-, -C(di-lower alkyl)-, -CH=CH- and -C(lower alkyl)=CH-; or R<sub>1</sub> together with an adjacent R<sub>2</sub>, or R<sub>1</sub> together with an adjacent R<sub>3</sub>, form a

-CH=CH- or a -CH=C(lower alkyl)- group;

u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sub>2</sub> is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R<sub>3</sub> is -CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, the R<sub>2</sub>'s can be the same or different; and provided that when u is 2 or 3, the R<sub>3</sub>'s can be the same or different;

R<sub>4</sub> is selected from B-(CH<sub>2</sub>)<sub>m</sub>C(O)-, wherein m is 0, 1, 2, 3, 4 or 5;

B-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 0, 1, 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein Z is -O-, -C(O)-, phenylene, -N(R<sub>8</sub>)- or -S(O)<sub>0-2</sub>-, e is 0, 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4, 5 or 6;

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-;

B-(C<sub>4</sub>-C<sub>6</sub> alkadienylene)-;

B-(CH<sub>2</sub>)<sub>t</sub>-Z-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1, 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0, 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6;


B-(CH<sub>2</sub>)<sub>t</sub>-V-(C<sub>2</sub>-C<sub>6</sub> alkenylene)- or

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-V-(CH<sub>2</sub>)<sub>t</sub>-, wherein V and t are as defined above, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6; B-(CH<sub>2</sub>)<sub>a</sub>-Z-(CH<sub>2</sub>)<sub>b</sub>-V-(CH<sub>2</sub>)<sub>d</sub>-, wherein Z and V are as defined above and a, b and d are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3, 4, 5 or 6; or


T-(CH<sub>2</sub>)<sub>s</sub>-, wherein T is cycloalkyl of 3-6 carbon atoms and s is 0, 1, 2, 3, 4, 5 or 6; or

R<sub>1</sub> and R<sub>4</sub> together form the group  $\text{B}-\overset{\text{I}}{\text{CH}=\text{C}}-$ ;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl, thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or



- W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy, alkoxyalkoxy, alkoxyalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl
- 5 lower alkanedioyl, allyloxy, -CF<sub>3</sub>, -OCF<sub>3</sub>, benzyl, R<sub>7</sub>-benzyl, benzyloxy, R<sub>7</sub>-benzyloxy, phenoxy, R<sub>7</sub>-phenoxy, dioxolanyl, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, OH, halogeno, -CN, -N<sub>3</sub>, -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, R<sub>11</sub>O<sub>2</sub>SNH-, (R<sub>11</sub>O<sub>2</sub>S)<sub>2</sub>N-, -S(O)<sub>2</sub>NH<sub>2</sub>, -S(O)<sub>0-2</sub>R<sub>8</sub>, tert-butyldimethyl-silyloxymethyl, -C(O)R<sub>12</sub>,
- 10 -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -CH=CHC(O)R<sub>12</sub>, -lower alkylene-C(O)R<sub>12</sub>, R<sub>10</sub>C(O)(lower alkyleneoxy)-, N(R<sub>8</sub>)(R<sub>9</sub>)C(O)(lower alkyleneoxy)- and



for substitution on ring carbon atoms,

- and the substituents on the substituted heteroaryl ring nitrogen atoms, when present, are selected from the group consisting of lower alkyl, lower alkoxy,
- 15 -C(O)OR<sub>10</sub>, -C(O)R<sub>10</sub>, OH, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, -S(O)<sub>2</sub>NH<sub>2</sub> and 2-(trimethylsilyl)-ethoxymethyl;


R<sub>7</sub> is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -COOH, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), OH, and halogeno;

R<sub>8</sub> and R<sub>9</sub> are independently selected from H or lower alkyl;

- 20 R<sub>10</sub> is selected from lower alkyl, phenyl, R<sub>7</sub>-phenyl, benzyl or R<sub>7</sub>-benzyl;

R<sub>11</sub> is selected from OH, lower alkyl, phenyl, benzyl, R<sub>7</sub>-phenyl or R<sub>7</sub>-benzyl;

R<sub>12</sub> is selected from H, OH, alkoxy, phenoxy, benzyloxy,



, -N(R<sub>8</sub>)(R<sub>9</sub>), lower alkyl, phenyl or R<sub>7</sub>-phenyl;


R<sub>13</sub> is selected from -O-, -CH<sub>2</sub>-, -NH-, -N(lower alkyl)- or -NC(O)R<sub>19</sub>;

- R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of H
- 25 and the groups defined for W; or R<sub>15</sub> is hydrogen and R<sub>16</sub> and R<sub>17</sub>, together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;

R<sub>19</sub> is H, lower alkyl, phenyl or phenyl lower alkyl; and


R20 and R21 are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzofused heteroaryl, W-substituted benzofused heteroaryl and cyclopropyl, wherein heteroaryl is as defined above.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (VIIA) or Formula (VIIB):



(VIIA)

or




(VIIB)


or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIIA) or Formula (VIIB) or of the isomers of Formula (VIIA) or Formula (VIIB), or prodrugs of the compounds of Formula (VIIA) or Formula (VIIB) or of the isomers, salts or solvates Formula (VIIA) or Formula (VIIB), wherein in Formula (VIIA) and Formula (VIIB):

A is  $-\text{CH}=\text{CH}-$ ,  $-\text{C}\equiv\text{C}-$  or  $-(\text{CH}_2)_p-$  wherein p is 0, 1 or 2;

B is



B' is




D is  $-(CH_2)_mC(O)-$  or  $-(CH_2)_q-$  wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;

E is  $C_{10}$  to  $C_{20}$  alkyl or  $-C(O)-(C_9 \text{ to } C_{19})\text{-alkyl}$ , wherein the alkyl is straight or branched, saturated or containing one or more double bonds;

5 R is hydrogen,  $C_1\text{-}C_{15}$  alkyl, straight or branched, saturated or containing one or more double bonds, or  $B-(CH_2)_r-$ , wherein r is 0, 1, 2, or 3;

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_1'$ ,  $R_2'$ , and  $R_3'$  are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino, dilower alkylamino,  $-NHC(O)OR_5$ ,  $R_6O_2SNH-$  and  $-S(O)_2NH_2$ ;

10  $R_4$  is




wherein n is 0, 1, 2 or 3;

$R_5$  is lower alkyl; and

$R_6$  is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents are 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino and dilower alkylamino.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (VIII):





(VIII)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIII) or of the isomers thereof, or prodrugs of the compounds of Formula (VIII) or of the isomers, salts or solvates thereof, wherein, in Formula (VIII) above,

$R_{26}$  is H or  $OG^1$ ;

G and G<sup>1</sup> are independently selected from the group consisting of



provided that when R<sup>26</sup> is H or

OH, G is not H;

R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy or -W-R<sup>30</sup>;

5 W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

10 R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

R<sup>30</sup> is selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, 15 R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, 20 imidazolyl and pyridyl;

R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy,




-CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyll or morpholinyll group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyll or morpholinyll group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone,

forms the spiro group ; and

R<sup>1</sup> is selected from the group consisting of


-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-E-(CH<sub>2</sub>)<sub>r</sub>-, wherein E is -O-, -C(O)-, phenylene, -NR<sup>22</sup>- or -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C<sub>2</sub>-C<sub>6</sub>)alkenylene-; and

-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R<sup>12</sup> is




R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl), -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;

provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1;

provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1;

provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and  
 provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;  
 and when Q is a bond, R<sup>1</sup> also can be:



5 M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub>)alkyl- and -C(di-(C<sub>1</sub>-C<sub>6</sub>)alkyl);

R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of  
 10 (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>,  
 -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>,  
 -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>,  
 -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>,  
 15 -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>15</sup> and R<sup>17</sup> are independently selected from the group consisting of  
 -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup> and -O(CO)NR<sup>19</sup>R<sup>20</sup>;

R<sup>16</sup> and R<sup>18</sup> are independently selected from the group consisting of H,  
 (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>15</sup> and R<sup>16</sup> together are =O, or R<sup>17</sup> and R<sup>18</sup> together are  
 20 =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

25 provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided  
 that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R<sup>1</sup> is 
$$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$$
, Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;


R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (IX) below:




or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (IX) or of the isomers thereof, or prodrugs of the compounds of Formula (IX) or of the isomers, salts or solvates thereof, wherein in Formula (IX):

R<sup>26</sup> is selected from the group consisting of:

- a) OH;
- b) OCH<sub>3</sub>;
- c) fluorine and
- d) chlorine.

R<sup>1</sup> is selected from the group consisting of



R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy and -W-R<sup>30</sup>;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

5 R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

R<sup>30</sup> is independently selected from the group consisting of R<sup>32</sup>-substituted T, 10 R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is independently selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

T is independently selected from the group consisting of phenyl, furyl, thienyl, 15 pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;


R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of H, halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>- 20 C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl,

piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

5 Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, or, with the 3-position ring carbon of the azetidinone,



R<sup>12</sup> is

10  $\begin{array}{c} | \\ -\text{CH}-, -\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})-, -\text{CF}-, -\text{C}(\text{OH})-, -\text{C}(\text{C}_6\text{H}_4\text{-R}^{23})-, -\text{N}-, \text{ or } -\text{NO}^+ \end{array}$  ;

R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl)-, -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

15 a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1; provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

20 R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>, -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>,  
25 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 August 2002 (01.08.2002)

PCT

(10) International Publication Number  
WO 02/058685 A2

(51) International Patent Classification<sup>7</sup>: A61K 31/00

(21) International Application Number: PCT/US02/02004

(22) International Filing Date: 25 January 2002 (25.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/264,275 26 January 2001 (26.01.2001) US  
60/323,842 21 September 2001 (21.09.2001) US

(71) Applicant: SCHERING CORPORATION [US/US];  
Patent Department - K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: DAVIS, Harry, R.; 64 River Bend Road,  
Berkeley Heights, NJ 07922 (US). KOSOGLU, Teddy;  
2457 Primrose Court, Jamison, Bucks County, PA  
18929-1178 (US).

(74) Agent: CANNONI, Ann, Marie; Schering-Plough Cor-  
poration, Patent Department K-6-1 1990, 2000 Galloping  
Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,  
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,  
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA,  
UZ, VN, YU, ZA, ZM.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: COMBINATIONS OF NICOTINIC ACID AND DERIVATIVES THEREOF AND STEROL ABSORPTION  
INHIBITOR(S) AND TREATMENTS FOR VASCULAR INDICATIONS

(57) Abstract: The present invention provides compositions, therapeutic combinations and methods including: (a) at least one of  
nicotinic acid or derivatives thereof; and (b) at least one substituted azetidinone or substituted b-lactam sterol absorption inhibitor  
which can be useful for treating vascular conditions, diabetes, obesity and lowering plasma levels of sterols.



WO 02/058685 A2

**COMBINATIONS OF NICOTINIC ACID AND DERIVATIVES THEREOF AND  
STEROL ABSORPTION INHIBITOR(S) AND TREATMENTS FOR VASCULAR  
INDICATIONS**

**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims the benefit of priority from U.S. Provisional Patent Application Serial No. 60/264,275 filed January 26, 2001 and U.S. Provisional Patent Application Serial No. 60/323,842 filed September 21, 2001, each of which is incorporated herein by reference.

**FIELD OF THE INVENTION**

The present invention relates to compositions and therapeutic combinations comprising nicotinic acid or derivatives thereof and certain sterol absorption inhibitors for treating hyperlipidemic conditions such as are associated with atherosclerosis, hypercholesterolemia and other vascular conditions in mammals.

**BACKGROUND OF THE INVENTION**

Atherosclerotic coronary heart disease (CHD) represents the major cause for death and vascular morbidity in the western world. Risk factors for atherosclerotic coronary heart disease include hypertension, diabetes mellitus, family history, male gender, cigarette smoke and serum cholesterol. A total cholesterol level in excess of 225-250 mg/dl is associated with significant elevation of risk of CHD.

Cholesteryl esters are a major component of atherosclerotic lesions and the major storage form of cholesterol in arterial wall cells. Formation of cholesteryl esters is also a key step in the intestinal absorption of dietary cholesterol. Thus, inhibition of cholesteryl ester formation and reduction of serum cholesterol can inhibit the progression of atherosclerotic lesion formation, decrease the accumulation of cholesteryl esters in the arterial wall, and block the intestinal absorption of dietary cholesterol.

The regulation of whole-body cholesterol homeostasis in mammals and animals involves the regulation of dietary cholesterol and modulation of cholesterol biosynthesis, bile acid biosynthesis and the catabolism of the cholesterol-containing plasma lipoproteins. The liver is the major organ responsible for cholesterol biosynthesis and catabolism and, for this reason, it is a prime determinant of plasma

cholesterol levels. The liver is the site of synthesis and secretion of very low density lipoproteins (VLDL) which are subsequently metabolized to low density lipoproteins (LDL) in the circulation. LDL are the predominant cholesterol-carrying lipoproteins in the plasma and an increase in their concentration is correlated with increased  
5 atherosclerosis. When intestinal cholesterol absorption is reduced, by whatever means, less cholesterol is delivered to the liver. The consequence of this action is decreased hepatic lipoprotein (VLDL) production and an increase in the hepatic clearance of plasma cholesterol, mostly as LDL. Thus, the net effect of inhibiting intestinal cholesterol absorption is a decrease in plasma cholesterol levels.

10 Nicotinic acid and its derivatives can inhibit hepatic production of VLDL and its metabolite LDL and increases HDL and apo A-1 levels.

U.S. Patents Nos. 5,767,115, 5,624,920, 5,668,990, 5,656,624 and 5,688,787, respectively, disclose hydroxy-substituted azetidinone compounds and substituted  $\beta$ -lactam compounds useful for lowering cholesterol and/or in inhibiting the formation of  
15 cholesterol-containing lesions in mammalian arterial walls. U.S. Patents Nos. 5,846,966 and 5,661,145, respectively, disclose hydroxy-substituted azetidinone compounds or substituted  $\beta$ -lactam compounds in combination with HMG CoA reductase inhibitors for preventing or treating atherosclerosis and reducing plasma cholesterol levels.

20 PCT Patent Application No. WO 00/38725 discloses cardiovascular therapeutic combinations including an ileal bile acid transport inhibitor or cholesteryl ester transport protein inhibitor in combination with a fibric acid derivative, nicotinic acid derivative, microsomal triglyceride transfer protein inhibitor, cholesterol absorption antagonist, phytosterol, stanol, antihypertensive agent or bile acid sequestrant.

25 U.S. Patent No. 5,698,527 discloses ergostanone derivatives substituted with disaccharides as cholesterol absorption inhibitors, employed alone or in combination with certain other cholesterol lowering agents, which are useful in the treatment of hypercholesterolemia and related disorders.


Despite recent improvements in the treatment of vascular disease, there  
30 remains a need in the art for improved compositions and treatments for hyperlipidaemia, atherosclerosis and other vascular conditions.

#### SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising:



(a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (I):



(I)

5 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (I) or of the isomers thereof, or prodrugs of the compounds of Formula (I) or of the isomers, salts or solvates thereof,

wherein in Formula (I) above:

Ar¹ and Ar² are independently selected from the group consisting of aryl and  
10 R⁴-substituted aryl;

Ar³ is aryl or R⁵-substituted aryl;

X, Y and Z are independently selected from the group consisting of  
-CH₂-, -CH(lower alkyl)- and -C(dilower alkyl)-;

R and R² are independently selected from the group consisting of -OR⁶, -  
15 O(CO)R⁶, -O(CO)OR⁹ and -O(CO)NR⁶R⁷;

R¹ and R³ are independently selected from the group consisting of hydrogen,  
lower alkyl and aryl;

q is 0 or 1;

r is 0 or 1;

20 m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;


R⁴ is 1-5 substituents independently selected from the group consisting of  
lower alkyl, -OR⁶, -O(CO)R⁶, -O(CO)OR⁹, -O(CH₂)₁-₅OR⁶, -O(CO)NR⁶R⁷,  
25 -NR⁶R⁷, -NR⁶(CO)R⁷, -NR⁶(CO)OR⁹, -NR⁶(CO)NR⁷R⁸, -NR⁶SO₂R⁹, -COOR⁶,  
-CONR⁶R⁷, -COR⁶, -SO₂NR⁶R⁷, S(O)₀-₂R⁹, -O(CH₂)₁-₁₀-COOR⁶,  
-O(CH₂)₁-₁₀CONR⁶R⁷, -(lower alkylene)COOR⁶, -CH=CH-COOR⁶, -CF₃, -CN,  
-NO₂ and halogen;

- $R^5$  is 1-5 substituents independently selected from the group consisting of -  
 $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
 $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ , -  
 $COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 5  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of  
 hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.


- In another embodiment, there is provided a composition comprising: (a) at least  
 10 one of nicotinic acid or derivatives thereof; and (b) a compound represented by  
 Formula (II) below:



(II)

- or pharmaceutically acceptable salt or solvate thereof, or prodrug of the compound of  
 15 Formula (II) or of the salt or solvate thereof.

In another embodiment, the present invention provides a composition  
 comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one  
 sterol absorption inhibitor represented by Formula (III):



(III)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (III) or of the isomers thereof, or prodrugs of the compounds of Formula (III) or of the isomers, salts or solvates thereof, wherein, in Formula (III) above:

$Ar^1$  is  $R^3$ -substituted aryl;  $Ar^2$  is  $R^4$ -substituted aryl;  $Ar^3$  is  $R^5$ -substituted aryl;

5 Y and Z are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(\text{lower alkyl})-$  and  $-C(\text{dilower alkyl})-$ ;

A is selected from  $-O-$ ,  $-S-$ ,  $-S(O)-$  or  $-S(O)_2-$ ;

$R^1$  is selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$  and  $-O(CO)NR^6R^7$ ;  $R^2$  is selected from the group consisting of hydrogen, lower alkyl and  
10 aryl; or  $R^1$  and  $R^2$  together are  $=O$ ;

q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;


$R^5$  is 1-3 substituents independently selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^9$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
15  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2\text{-lower alkyl}$ ,  $-NR^6SO_2\text{-aryl}$ ,  $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}\text{-alkyl}$ ,  $S(O)_{0-2}\text{-aryl}$ ,  $-O(CH_2)_{1-10}\text{-COOR}^6$ ,  $-O(CH_2)_{1-10}\text{CONR}^6R^7$ , o-halogeno, m-halogeno, o-lower alkyl, m-lower alkyl,  $-(\text{lower alkylene})\text{-COOR}^6$ , and  $-\text{CH}=\text{CH}\text{-COOR}^6$ ;

$R^3$  and  $R^4$  are independently 1-3 substituents independently selected from the  
20 group consisting of  $R^5$ , hydrogen, p-lower alkyl, aryl,  $-\text{NO}_2$ ,  $-\text{CF}_3$  and p-halogeno;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.

25 In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (IV):



(IV)


or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 5 of Formula (IV) or of the isomers thereof, or prodrugs of the compounds of Formula  
 (IV) or of the isomers, salts or solvates thereof, wherein, in Formula (IV) above:

A is selected from the group consisting of  $R^2$ -substituted heterocycloalkyl,  $R^2$ -  
 substituted heteroaryl,  $R^2$ -substituted benzofused heterocycloalkyl, and  $R^2$ -substituted  
 benzofused heteroaryl;

10  $Ar^1$  is aryl or  $R^3$ -substituted aryl;

$Ar^2$  is aryl or  $R^4$ -substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the



$R^1$  is selected from the group consisting of:


15  $-(CH_2)_q-$ , wherein q is 2-6, provided that when Q forms a spiro ring, q can  
 also be zero or 1;

$-(CH_2)_e-G-(CH_2)_r-$ , wherein G is  $-O-$ ,  $-C(O)-$ , phenylene,  $-NR^8-$  or  
 $-S(O)_{0-2}-$ , e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

$-(C_2-C_6 \text{ alkenylene})-$ ; and

20  $-(CH_2)_fV-(CH_2)_g-$ , wherein V is  $C_3-C_6$  cycloalkylene, f is 1-5 and g is 0-5,  
 provided that the sum of f and g is 1-6;

$R^5$  is selected from:




$R^6$  and  $R^7$  are independently selected from the group consisting of  
 25  $-CH_2-$ ,  $-CH(C_1-C_6 \text{ alkyl})-$ ,  $-C(di-(C_1-C_6 \text{ alkyl}))-$ ,  $-CH=CH-$  and

$-\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})=\text{CH}-$ ; or  $\text{R}^5$  together with an adjacent  $\text{R}^6$ , or  $\text{R}^5$  together with an adjacent  $\text{R}^7$ , form a  $-\text{CH}=\text{CH}-$  or a  $-\text{CH}=\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})-$  group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when  $\text{R}^6$  is  $-\text{CH}=\text{CH}-$  or  $-\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})=\text{CH}-$ , a is 1; provided that when  $\text{R}^7$  is  $-\text{CH}=\text{CH}-$  or  $-\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})=\text{CH}-$ , b is 1; provided that when a is 2 or 3, the  $\text{R}^6$ 's can be the same or different; and provided that when b is 2 or 3, the  $\text{R}^7$ 's can be the same or different;

and when Q is a bond,  $\text{R}^1$  also can be selected from:



where M is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O})-$  or  $-\text{S}(\text{O})_2-$ ;

X, Y and Z are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{CH}(\text{C}_1\text{-C}_6 \text{ alkyl})-$  and  $-\text{C}(\text{di}(\text{C}_1\text{-C}_6 \text{ alkyl}))-$ ;

$\text{R}^{10}$  and  $\text{R}^{12}$  are independently selected from the group consisting of  $-\text{OR}^{14}$ ,  $-\text{O}(\text{CO})\text{R}^{14}$ ,  $-\text{O}(\text{CO})\text{OR}^{16}$  and  $-\text{O}(\text{CO})\text{NR}^{14}\text{R}^{15}$ ;

$\text{R}^{11}$  and  $\text{R}^{13}$  are independently selected from the group consisting of hydrogen,  $(\text{C}_1\text{-C}_6)\text{alkyl}$  and aryl; or  $\text{R}^{10}$  and  $\text{R}^{11}$  together are  $=\text{O}$ , or  $\text{R}^{12}$  and  $\text{R}^{13}$  together are  $=\text{O}$ ;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;


s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;


v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

$\text{R}^2$  is 1-3 substituents on the ring carbon atoms selected from the group consisting of hydrogen,  $(\text{C}_1\text{-C}_{10})\text{alkyl}$ ,  $(\text{C}_2\text{-C}_{10})\text{alkenyl}$ ,  $(\text{C}_2\text{-C}_{10})\text{alkynyl}$ ,  $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ ,  $(\text{C}_3\text{-C}_6)\text{cycloalkenyl}$ ,  $\text{R}^{17}$ -substituted aryl,  $\text{R}^{17}$ -substituted benzyl,  $\text{R}^{17}$ -substituted benzyloxy,  $\text{R}^{17}$ -substituted aryloxy, halogeno,  $-\text{NR}^{14}\text{R}^{15}$ ,  $\text{NR}^{14}\text{R}^{15}(\text{C}_1\text{-C}_6 \text{ alkylene})$ ,  $\text{NR}^{14}\text{R}^{15}\text{C}(\text{O})(\text{C}_1\text{-C}_6 \text{ alkylene})$ ,  $-\text{NHC}(\text{O})\text{R}^{16}$ ,  $\text{OH}$ ,

$C_1-C_6$  alkoxy,  $-OC(O)R^{16}$ ,  $-COR^{14}$ , hydroxy( $C_1-C_6$ )alkyl,  
 $(C_1-C_6)$ alkoxy( $C_1-C_6$ )alkyl,  $NO_2$ ,  $-S(O)_{0-2}R^{16}$ ,  $-SO_2NR^{14}R^{15}$  and  
 $-(C_1-C_6 \text{ alkylene})COOR^{14}$ ; when  $R^2$  is a substituent on a heterocycloalkyl ring,  $R^2$  is

as defined, or is  $=O$  or ; and, where  $R^2$  is a substituent on a  
 5 substitutable ring nitrogen, it is hydrogen, ( $C_1-C_6$ )alkyl, aryl, ( $C_1-C_6$ )alkoxy, aryloxy,  
 $(C_1-C_6)$ alkylcarbonyl, arylcarbonyl, hydroxy,  $-(CH_2)_{1-6}CONR^{18}R^{18}$ ,



wherein J is  $-O-$ ,  $-NH-$ ,  $-NR^{18}-$  or  $-CH_2-$ ;

$R^3$  and  $R^4$  are independently selected from the group consisting of 1-3  
 10 substituents independently selected from the group consisting of ( $C_1-C_6$ )alkyl,  
 $-OR^{14}$ ,  $-O(CO)R^{14}$ ,  $-O(CO)OR^{16}$ ,  $-O(CH_2)_{1-5}OR^{14}$ ,  $-O(CO)NR^{14}R^{15}$ ,  $-NR^{14}R^{15}$ ,  
 $-NR^{14}(CO)R^{15}$ ,  $-NR^{14}(CO)OR^{16}$ ,  $-NR^{14}(CO)NR^{15}R^{19}$ ,  $-NR^{14}SO_2R^{16}$ ,  $-COOR^{14}$ ,  
 $-CONR^{14}R^{15}$ ,  $-COR^{14}$ ,  $-SO_2NR^{14}R^{15}$ ,  $S(O)_{0-2}R^{16}$ ,  $-O(CH_2)_{1-10}-COOR^{14}$ ,  
 $-O(CH_2)_{1-10}CONR^{14}R^{15}$ ,  $-(C_1-C_6 \text{ alkylene})-COOR^{14}$ ,  $-CH=CH-COOR^{14}$ ,  $-CF_3$ ,  $-CN$ ,  $-$   
 15  $NO_2$  and halogen;

$R^8$  is hydrogen, ( $C_1-C_6$ )alkyl, aryl ( $C_1-C_6$ )alkyl,  $-C(O)R^{14}$  or  $-COOR^{14}$ ;

$R^9$  and  $R^{17}$  are independently 1-3 groups independently selected from the  
 group consisting of hydrogen, ( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxy,  $-COOH$ ,  $NO_2$ ,  
 $-NR^{14}R^{15}$ ,  $OH$  and halogeno;


20  $R^{14}$  and  $R^{15}$  are independently selected from the group consisting of hydrogen,  
 $(C_1-C_6)$ alkyl, aryl and aryl-substituted ( $C_1-C_6$ )alkyl;

$R^{16}$  is ( $C_1-C_6$ )alkyl, aryl or  $R^{17}$ -substituted aryl;

$R^{18}$  is hydrogen or ( $C_1-C_6$ )alkyl; and

$R^{19}$  is hydrogen, hydroxy or ( $C_1-C_6$ )alkoxy.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (V):



(V)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (V) or of the isomers thereof, or prodrugs of the compounds of Formula (V) or of the isomers, salts or solvates thereof, wherein, in Formula (V) above:

- 10  $Ar^1$  is aryl,  $R^{10}$ -substituted aryl or heteroaryl;  
 $Ar^2$  is aryl or  $R^4$ -substituted aryl;  
 $Ar^3$  is aryl or  $R^5$ -substituted aryl;  
X and Y are independently selected from the group consisting of  $-CH_2-$ ,  
 $-CH(\text{lower alkyl})-$  and  $-C(\text{dilower alkyl})-$ ;  
15 R is  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$  or  $-O(CO)NR^6R^7$ ;  $R^1$  is hydrogen, lower alkyl or aryl; or R and  $R^1$  together are  $=O$ ;  
q is 0 or 1;  
r is 0, 1 or 2;  
m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and  
20 q is 1, 2, 3, 4 or 5;  
 $R^4$  is 1-5 substituents independently selected from the group consisting of  
lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  
 $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  
 $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
25  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;  
 $R^5$  is 1-5 substituents independently selected from the group consisting of -  
 $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,

-NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, halogen, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;


R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of  
 5 hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;

R<sup>9</sup> is lower alkyl, aryl or aryl-substituted lower alkyl; and

R<sup>10</sup> is 1-5 substituents independently selected from the group consisting of  
 lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>,  
 -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>,  
 10 -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen.

In another embodiment, the present invention provides a composition  
 comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one  
 sterol absorption inhibitor represented by Formula (VI):

15



(VI)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 of Formula (VI) or of the isomers thereof, or prodrugs of the compounds of Formula  
 20 (VI) or of the isomers, salts or solvates thereof, wherein:

R<sub>1</sub> is

$\overset{|}{-}\text{CH}-$ ,  $\overset{|}{-}\text{C}(\text{lower alkyl})-$ ,  $\overset{|}{-}\text{CF}-$ ,  $\overset{|}{-}\text{C}(\text{OH})-$ ,  $\overset{|}{-}\text{C}(\text{C}_6\text{H}_5)-$ ,  $\overset{|}{-}\text{C}(\text{C}_6\text{H}_4-\text{R}_{15})-$ ,  
 $\overset{|}{-}\text{N}-$  or  $\overset{|}{-}\text{N}^+\text{O}^-$ ;

R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of:  
 25 -CH<sub>2</sub>-, -CH(lower alkyl)-, -C(di-lower alkyl)-, -CH=CH- and -C(lower alkyl)=CH-; or  
 R<sub>1</sub> together with an adjacent R<sub>2</sub>, or R<sub>1</sub> together with an adjacent R<sub>3</sub>, form a



-CH=CH- or a -CH=C(lower alkyl)- group;

u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sub>2</sub> is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R<sub>3</sub> is -CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, the R<sub>2</sub>'s can be the same or different; and provided that when u is 2 or 3, the R<sub>3</sub>'s can be the same or different;

R<sub>4</sub> is selected from B-(CH<sub>2</sub>)<sub>m</sub>C(O)-, wherein m is 0, 1, 2, 3, 4 or 5;

B-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 0, 1, 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein Z is -O-, -C(O)-, phenylene, -N(R<sub>8</sub>)- or -S(O)<sub>0-2</sub>-, e is 0, 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4, 5 or 6;

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-;

B-(C<sub>4</sub>-C<sub>6</sub> alkadienylene)-;

B-(CH<sub>2</sub>)<sub>t</sub>-Z-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1, 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0, 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6;


B-(CH<sub>2</sub>)<sub>t</sub>-V-(C<sub>2</sub>-C<sub>6</sub> alkenylene)- or

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-V-(CH<sub>2</sub>)<sub>t</sub>-, wherein V and t are as defined above, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6; B-(CH<sub>2</sub>)<sub>a</sub>-Z-(CH<sub>2</sub>)<sub>b</sub>-V-(CH<sub>2</sub>)<sub>d</sub>-, wherein Z and V are as defined above and a, b and d are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3, 4, 5 or 6; or


T-(CH<sub>2</sub>)<sub>s</sub>-, wherein T is cycloalkyl of 3-6 carbon atoms and s is 0, 1, 2, 3, 4, 5 or 6; or

R<sub>1</sub> and R<sub>4</sub> together form the group  $\text{B}-\text{CH}=\overset{\text{I}}{\text{C}}-$ ;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl, thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or



W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy, alkoxycarbonylalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl  
 5 lower alkanedioyl, allyloxy, -CF<sub>3</sub>, -OCF<sub>3</sub>, benzyl, R<sub>7</sub>-benzyl, benzyloxy, R<sub>7</sub>-benzyloxy, phenoxy, R<sub>7</sub>-phenoxy, dioxolanyl, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, OH, halogeno, -CN, -N<sub>3</sub>, -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, R<sub>11</sub>O<sub>2</sub>SNH-, (R<sub>11</sub>O<sub>2</sub>S)<sub>2</sub>N-, -S(O)<sub>2</sub>NH<sub>2</sub>, -S(O)<sub>0-2</sub>R<sub>8</sub>, tert-butyldimethyl-silyloxymethyl, -C(O)R<sub>12</sub>,  
 10 -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -CH=CHC(O)R<sub>12</sub>, -lower alkylene-C(O)R<sub>12</sub>, R<sub>10</sub>C(O)(lower alkyleneoxy)-, N(R<sub>8</sub>)(R<sub>9</sub>)C(O)(lower alkyleneoxy)- and



for substitution on ring carbon atoms,

and the substituents on the substituted heteroaryl ring nitrogen atoms, when present, are selected from the group consisting of lower alkyl, lower alkoxy,  
 15 -C(O)OR<sub>10</sub>, -C(O)R<sub>10</sub>, OH, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, -S(O)<sub>2</sub>NH<sub>2</sub> and 2-(trimethylsilyl)-ethoxymethyl;


R<sub>7</sub> is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -COOH, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), OH, and halogeno;

R<sub>8</sub> and R<sub>9</sub> are independently selected from H or lower alkyl;

20 R<sub>10</sub> is selected from lower alkyl, phenyl, R<sub>7</sub>-phenyl, benzyl or R<sub>7</sub>-benzyl;

R<sub>11</sub> is selected from OH, lower alkyl, phenyl, benzyl, R<sub>7</sub>-phenyl or R<sub>7</sub>-benzyl;

R<sub>12</sub> is selected from H, OH, alkoxy, phenoxy, benzyloxy,



, -N(R<sub>8</sub>)(R<sub>9</sub>), lower alkyl, phenyl or R<sub>7</sub>-phenyl;


R<sub>13</sub> is selected from -O-, -CH<sub>2</sub>-, -NH-, -N(lower alkyl)- or -NC(O)R<sub>19</sub>;

R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of H  
 25 and the groups defined for W; or R<sub>15</sub> is hydrogen and R<sub>16</sub> and R<sub>17</sub>, together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;

R<sub>19</sub> is H, lower alkyl, phenyl or phenyl lower alkyl; and


R<sub>20</sub> and R<sub>21</sub> are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzofused heteroaryl, W-substituted benzofused heteroaryl and cyclopropyl, wherein heteroaryl is as defined above.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (VIIA) or Formula (VIIB):



(VIIA)

or




(VIIB)


or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIIA) or Formula (VIIB) or of the isomers of Formula (VIIA) or Formula (VIIB), or prodrugs of the compounds of Formula (VIIA) or Formula (VIIB) or of the isomers, salts or solvates Formula (VIIA) or Formula (VIIB), wherein in Formula (VIIA) and Formula (VIIB):

A is  $-\text{CH}=\text{CH}-$ ,  $-\text{C}\equiv\text{C}-$  or  $-(\text{CH}_2)_p-$  wherein p is 0, 1 or 2;

B is



B' is




D is  $-(CH_2)_mC(O)-$  or  $-(CH_2)_q-$  wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;

E is  $C_{10}$  to  $C_{20}$  alkyl or  $-C(O)-(C_9 \text{ to } C_{19})\text{-alkyl}$ , wherein the alkyl is straight or branched, saturated or containing one or more double bonds;

5 R is hydrogen,  $C_1$ - $C_{15}$  alkyl, straight or branched, saturated or containing one or more double bonds, or  $B-(CH_2)_r-$ , wherein r is 0, 1, 2, or 3;

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_1'$ ,  $R_2'$ , and  $R_3'$  are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino, dilower alkylamino,  $-NHC(O)OR_5$ ,  $R_6O_2SNH-$  and  $-S(O)_2NH_2$ ;

10  $R_4$  is




wherein n is 0, 1, 2 or 3;

$R_5$  is lower alkyl; and

15  $R_6$  is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents are 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino and dilower alkylamino.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (VIII):

20





(VIII)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIII) or of the isomers thereof, or prodrugs of the compounds of Formula (VIII) or of the isomers, salts or solvates thereof, wherein, in Formula (VIII) above,

$R^{26}$  is H or  $OG^1$ ;

G and G<sup>1</sup> are independently selected from the group consisting of



provided that when R<sup>26</sup> is H or

OH, G is not H;

R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy or -W-R<sup>30</sup>;

5 W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

10 R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

R<sup>30</sup> is selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, 15 R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, 20 imidazolyl and pyridyl;


R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy,

-CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indoliny or morpholiny group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indoliny or morpholiny group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone,

forms the spiro group ; and

R<sup>1</sup> is selected from the group consisting of


-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-E-(CH<sub>2</sub>)<sub>r</sub>-, wherein E is -O-, -C(O)-, phenylene, -NR<sup>22</sup>- or -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C<sub>2</sub>-C<sub>6</sub>)alkenylene-; and

-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R<sup>12</sup> is




R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl), -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;

provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1;

provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1;

provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and  
 provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;  
 and when Q is a bond, R<sup>1</sup> also can be:



5 M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub>)alkyl- and -C(di-(C<sub>1</sub>-C<sub>6</sub>)alkyl);

R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3  
 substituents independently selected from the group consisting of  
 10 (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>,  
 -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>,  
 -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>,  
 -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>,  
 15 -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>15</sup> and R<sup>17</sup> are independently selected from the group consisting of  
 -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup> and -O(CO)NR<sup>19</sup>R<sup>20</sup>;

R<sup>16</sup> and R<sup>18</sup> are independently selected from the group consisting of H,  
 (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>15</sup> and R<sup>16</sup> together are =O, or R<sup>17</sup> and R<sup>18</sup> together are  
 20 =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

25 provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided  
 that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R<sup>1</sup> is 
$$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$$
, Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;


R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

In another embodiment, the present invention provides a composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one sterol absorption inhibitor represented by Formula (IX) below:




or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (IX) or of the isomers thereof, or prodrugs of the compounds of Formula (IX) or of the isomers, salts or solvates thereof, wherein in Formula (IX):

R<sup>26</sup> is selected from the group consisting of:

- a) OH;
- b) OCH<sub>3</sub>;
- c) fluorine and
- d) chlorine.

R<sup>1</sup> is selected from the group consisting of





R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy and -W-R<sup>30</sup>;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

5        R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

10        R<sup>30</sup> is independently selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is independently selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

15        T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;


20        R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of H, halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl,

piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

5 Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, or, with the 3-position ring carbon of the azetidinone,



R<sup>12</sup> is

10  $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$ ,  $\begin{array}{c} | \\ -\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})- \\ | \end{array}$ ,  $\begin{array}{c} | \\ -\text{CF}- \\ | \end{array}$ ,  $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$ ,  $\begin{array}{c} | \\ -\text{C}(\text{C}_6\text{H}_4\text{-R}^{23})- \\ | \end{array}$ ,  $\begin{array}{c} | \\ -\text{N}- \\ | \end{array}$ , or  $\begin{array}{c} | \\ -\text{N}^+\text{O}^- \\ | \end{array}$  ;

R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub> alkyl))- , -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

15 a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1; provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

20 R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>, -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>,  
25 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

5 R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

10 R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

In another embodiment, a composition is provided which comprises: (a) at least one of nicotinic acid or its derivatives; and (b) at least one substituted azetidinone compound or at least one substituted  $\beta$ -lactam compound, or isomers of the at least one substituted azetidinone compound or of the at least one substituted  $\beta$ -lactam  
15 compound, or pharmaceutically acceptable salts or solvates of the at least one substituted azetidinone compound or of the at least one substituted  $\beta$ -lactam compound or of the isomers thereof, or prodrugs of the at least one substituted azetidinone compound or of the at least one substituted  $\beta$ -lactam compound or of the isomers, salts or solvates thereof.

20 Therapeutic combinations are provided comprising: (a) a first amount of at least one of nicotinic acid or derivatives thereof; and (b) a second amount of at least one sterol absorption inhibitor represented by Formulae (I-XI), substituted azetidinone compounds, substituted  $\beta$ -lactam compounds, isomers of the sterol absorption inhibitors, or pharmaceutically acceptable salts or solvates of the sterol absorption  
25 inhibitors, or prodrugs of the sterol absorption inhibitors or of the isomers, salts or solvates thereof, wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

30 Pharmaceutical compositions for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a

mammal, comprising a therapeutically effective amount of the above compositions or therapeutic combinations and a pharmaceutically acceptable carrier also are provided.

Methods of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of  
5 administering to a mammal in need of such treatment an effective amount of the above compositions or therapeutic combinations also are provided.

Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by  
10 the term "about."

### **DETAILED DESCRIPTION**

In one embodiment, the present invention is directed to compositions, pharmaceutical compositions, therapeutic combinations, kits and methods of  
15 treatment using the same comprising at least one (one or more) of nicotinic acid (niacin) or derivatives thereof(s) and at least one (one or more) of substituted azetidinone sterol absorption inhibitors or substituted  $\beta$ -lactam sterol absorption inhibitors such as are discussed in detail below.

As used herein, "nicotinic acid derivative" means a compound comprising a  
20 pyridine-3-carboxylate structure or a pyrazine-2-carboxylate structure, including acid forms, salts, esters, zwitterions and tautomers, where available. Examples of nicotinic acid derivatives include niceritrol, nicofuranose and acipimox (5-methyl pyrazine-2-carboxylic acid 4-oxide). Nicotinic acid and its derivatives inhibit hepatic production of VLDL and its metabolite LDL and increases HDL and apo A-1 levels. An example of a  
25 suitable nicotinic acid product is NIASPAN® (niacin extended-release tablets) which are available from Kos.

The nicotinic acid or derivatives thereof(s) are administered in a therapeutically effective amount to treat the specified condition, for example a total daily dosage of nicotinic acid or a derivative thereof can range from about 500 to about 10,000  
30 mg/day, preferably about 1000 to about 8000 mg/day, and more preferably about 3000 to about 6000 mg/day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on such factors as the potency of the compound administered, the age, weight, condition and response of the patient.


The phrase "therapeutically effective amount" means that amount of a therapeutic agent of the composition, such as the nicotinic acid or derivatives thereof(s), sterol absorption inhibitor(s) and pharmacological or therapeutic agents described below, that will elicit a biological or medical response of a tissue, system, animal or mammal that is being sought by the administrator (such as a researcher, doctor or veterinarian) which includes alleviation of the symptoms of the condition or disease being treated and the prevention, slowing or halting of progression of one or more conditions, for example vascular conditions, such as hyperlipidemia (for example atherosclerosis, hypercholesterolemia or sitosterolemia), vascular inflammation, stroke, diabetes, obesity and/or to reduce the level of sterol(s) (such as cholesterol) in the plasma.

As used herein, "combination therapy" or "therapeutic combination" means the administration of two or more therapeutic agents, such as nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s), to prevent or treat a condition, for example a vascular condition, such as hyperlipidaemia (for example atherosclerosis, hypercholesterolemia or sitosterolemia), stroke, diabetes, obesity and/or reduce the level of sterol(s) in the plasma. As used herein, "vascular" comprises cardiovascular, cerebrovascular and combinations thereof. Such administration includes coadministration of these therapeutic agents in a substantially simultaneous manner, such as in a single tablet or capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each therapeutic agent. Also, such administration includes use of each type of therapeutic agent in a sequential manner. In either case, the treatment using the combination therapy will provide beneficial effects in treating the condition. A potential advantage of the combination therapy disclosed herein may be a reduction in the required amount of an individual therapeutic compound or the overall total amount of therapeutic compounds that are effective in treating the condition. By using a combination of therapeutic agents, the side effects of the individual compounds can be reduced as compared to a monotherapy, which can improve patient compliance. Also, therapeutic agents can be selected to provide a broader range of complementary effects or complimentary modes of action.

As discussed above, the compositions, pharmaceutical compositions and therapeutic combinations of the present invention comprise one or more substituted azetidinone or substituted  $\beta$ -lactam sterol absorption inhibitors discussed in detail below. As used herein, "sterol absorption inhibitor" means a compound capable of

inhibiting the absorption of one or more sterols, including but not limited to cholesterol, phytosterols (such as sitosterol, campesterol, stigmasterol and avenosterol), 5 $\alpha$ -stanols (such as cholestanol, 5 $\alpha$ -campestanol, 5 $\alpha$ -sitostanol), and mixtures thereof, when administered in a therapeutically effective (sterol absorption inhibiting) amount to a mammal.

In a preferred embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (I) below:



(I)

or isomers of the compounds of Formula (I), or pharmaceutically acceptable salts or solvates of the compounds of Formula (I) or of the isomers of the compounds of Formula (I), or prodrugs of the compounds of Formula (I) or of the isomers, salts or solvates of the compounds of Formula (I), wherein, in Formula (I) above:

$\text{Ar}^1$  and  $\text{Ar}^2$  are independently selected from the group consisting of aryl and  $\text{R}^4$ -substituted aryl;

$\text{Ar}^3$  is aryl or  $\text{R}^5$ -substituted aryl;

X, Y and Z are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{CH}(\text{lower alkyl})-$  and  $-\text{C}(\text{dilower alkyl})-$ ;

R and  $\text{R}^2$  are independently selected from the group consisting of  $-\text{OR}^6$ ,  $-\text{O}(\text{CO})\text{R}^6$ ,  $-\text{O}(\text{CO})\text{OR}^9$  and  $-\text{O}(\text{CO})\text{NR}^6\text{R}^7$ ;

$\text{R}^1$  and  $\text{R}^3$  are independently selected from the group consisting of hydrogen, lower alkyl and aryl;

q is 0 or 1; r is 0 or 1; m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3,

4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

$R^4$  is 1-5 substituents independently selected from the group consisting of lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  
 5  $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  
 $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$ ,  $-CH=CH-COOR^6$ ,  $-CF_3$ ,  $-CN$ ,  
 $-NO_2$  and halogen;

$R^5$  is 1-5 substituents independently selected from the group consisting of -  
 10  $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
 $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ ,  
 $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of  
 15 hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.

Preferably,  $R^4$  is 1-3 independently selected substituents, and  $R^5$  is preferably 1-3 independently selected substituents.

As used herein, the term "alkyl" or "lower alkyl" means straight or branched  
 20 alkyl chains having from 1 to 6 carbon atoms and "alkoxy" means alkoxy groups having 1 to 6 carbon atoms. Non-limiting examples of lower alkyl groups include, for example methyl, ethyl, propyl, and butyl groups.

"Alkenyl" means straight or branched carbon chains having one or more double bonds in the chain, conjugated or unconjugated. Similarly, "alkynyl" means straight or  
 25 branched carbon chains having one or more triple bonds in the chain. Where an alkyl, alkenyl or alkynyl chain joins two other variables and is therefore bivalent, the terms alkylene, alkenylene and alkynylene are used.

"Cycloalkyl" means a saturated carbon ring of 3 to 6 carbon atoms, while "cycloalkylene" refers to a corresponding bivalent ring, wherein the points of  
 30 attachment to other groups include all positional isomers.

"Halogeno" refers to fluorine, chlorine, bromine or iodine radicals.

"Aryl" means phenyl, naphthyl, indenyl, tetrahydronaphthyl or indanyl.

"Phenylene" means a bivalent phenyl group, including ortho, meta and para-substitution.

The statements wherein, for example, R, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup>, are said to be  
5 independently selected from a group of substituents, mean that R, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are  
independently selected, but also that where an R, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> variable occurs more  
than once in a molecule, each occurrence is independently selected (e.g., if R is -OR<sup>6</sup>,  
wherein R<sup>6</sup> is hydrogen, R<sup>2</sup> can be -OR<sup>6</sup> wherein R<sup>6</sup> is lower alkyl). Those skilled in  
the art will recognize that the size and nature of the substituent(s) will affect the  
10 number of substituents that can be present.

Compounds of the invention have at least one asymmetrical carbon atom and  
therefore all isomers, including enantiomers, stereoisomers, rotamers, tautomers and  
racemates of the compounds of Formulae (I-XI) (where they exist) are contemplated  
as being part of this invention. The invention includes d and l isomers in both pure  
15 form and in admixture, including racemic mixtures. Isomers can be prepared using  
conventional techniques, either by reacting optically pure or optically enriched starting  
materials or by separating isomers of a compound of the Formulae I-XI. Isomers may  
also include geometric isomers, e.g., when a double bond is present..

Those skilled in the art will appreciate that for some of the compounds of the  
20 Formulas I-XI, one isomer will show greater pharmacological activity than other  
isomers.

Compounds of the invention with an amino group can form pharmaceutically  
acceptable salts with organic and inorganic acids. Examples of suitable acids for salt  
formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic,  
25 malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and  
carboxylic acids well known to those in the art. The salt is prepared by contacting the  
free base form with a sufficient amount of the desired acid to produce a salt. The free  
base form may be regenerated by treating the salt with a suitable dilute aqueous base  
solution such as dilute aqueous sodium bicarbonate. The free base form differs from  
30 its respective salt form somewhat in certain physical properties, such as solubility in  
polar solvents, but the salt is otherwise equivalent to its respective free base forms for  
purposes of the invention.



Certain compounds of the invention are acidic (e.g., those compounds which possess a carboxyl group). These compounds form pharmaceutically acceptable salts with inorganic and organic bases. Examples of such salts are the sodium, potassium, calcium, aluminum, gold and silver salts. Also included are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.

As used herein, "solvate" means a molecular or ionic complex of molecules or ions of solvent with those of solute (for example, one or more compounds of Formulae I-XI, isomers of the compounds of Formulae I-XI, or prodrugs of the compounds of Formulae I-XI). Non-limiting examples of useful solvents include polar, protic solvents such as water and/or alcohols (for example methanol).

As used herein, "prodrug" means compounds that are drug precursors which, following administration to a patient, release the drug *in vivo* via some chemical or physiological process (e.g., a prodrug on being brought to the physiological pH or through enzyme action is converted to the desired drug form).

Preferred compounds of Formula (I) are those in which  $Ar^1$  is phenyl or  $R^4$ -substituted phenyl, more preferably (4- $R^4$ )-substituted phenyl.  $Ar^2$  is preferably phenyl or  $R^4$ -substituted phenyl, more preferably (4- $R^4$ )-substituted phenyl.  $Ar^3$  is preferably  $R^5$ -substituted phenyl, more preferably (4- $R^5$ )-substituted phenyl. When  $Ar^1$  is (4- $R^4$ )-substituted phenyl,  $R^4$  is preferably a halogen. When  $Ar^2$  and  $Ar^3$  are  $R^4$ - and  $R^5$ -substituted phenyl, respectively,  $R^4$  is preferably halogen or  $-OR^6$  and  $R^5$  is preferably  $-OR^6$ , wherein  $R^6$  is lower alkyl or hydrogen. Especially preferred are compounds wherein each of  $Ar^1$  and  $Ar^2$  is 4-fluorophenyl and  $Ar^3$  is 4-hydroxyphenyl or 4-methoxyphenyl.

X, Y and Z are each preferably  $-CH_2-$ .  $R^1$  and  $R^3$  are each preferably hydrogen.  $R$  and  $R^2$  are preferably  $-OR^6$  wherein  $R^6$  is hydrogen, or a group readily metabolizable to a hydroxyl (such as  $-O(CO)R^6$ ,  $-O(CO)OR^9$  and  $-O(CO)NR^6R^7$ , defined above).


The sum of m, n, p, q and r is preferably 2, 3 or 4, more preferably 3. Preferred are compounds wherein m, n and r are each zero, q is 1 and p is 2.

Also preferred are compounds of Formula (I) in which p, q and n are each zero, r is 1 and m is 2 or 3. More preferred are compounds wherein m, n and r are each zero, q is 1, p is 2, Z is  $-\text{CH}_2-$  and R is  $-\text{OR}^6$ , especially when  $\text{R}^6$  is hydrogen.

Also more preferred are compounds of Formula (I) wherein p, q and n are each zero, r is 1, m is 2, X is  $-\text{CH}_2-$  and  $\text{R}^2$  is  $-\text{OR}^6$ , especially when  $\text{R}^6$  is hydrogen.

Another group of preferred compounds of Formula (I) is that in which  $\text{Ar}^1$  is phenyl or  $\text{R}^4$ -substituted phenyl,  $\text{Ar}^2$  is phenyl or  $\text{R}^4$ -substituted phenyl and  $\text{Ar}^3$  is  $\text{R}^5$ -substituted phenyl. Also preferred are compounds in which  $\text{Ar}^1$  is phenyl or  $\text{R}^4$ -substituted phenyl,  $\text{Ar}^2$  is phenyl or  $\text{R}^4$ -substituted phenyl,  $\text{Ar}^3$  is  $\text{R}^5$ -substituted phenyl, and the sum of m, n, p, q and r is 2, 3 or 4, more preferably 3. More preferred are compounds wherein  $\text{Ar}^1$  is phenyl or  $\text{R}^4$ -substituted phenyl,  $\text{Ar}^2$  is phenyl or  $\text{R}^4$ -substituted phenyl,  $\text{Ar}^3$  is  $\text{R}^5$ -substituted phenyl, and wherein m, n and r are each zero, q is 1 and p is 2, or wherein p, q and n are each zero, r is 1 and m is 2 or 3.

In a preferred embodiment, a sterol inhibitor of Formula (I) (ezetimibe) useful in the compositions, therapeutic combinations and methods of the present invention is represented by Formula (II) below:




(II)

or pharmaceutically acceptable salts or solvates of the compound of Formula (II), or prodrugs of the compound of Formula (II) or of the salts or solvates of the compound of Formula (II).

Compounds of Formula I can be prepared by a variety of methods well known to those skilled in the art, for example such as are disclosed in U.S. Patents Nos. 5,631,365, 5,767,115, 5,846,966, 6,207,822 and U.S. Provisional Patent Application No. 60/279,288 filed March 28, 2001, and PCT Patent Application WO 93/02048, each of which is incorporated herein by reference, and in the Example below. For example,


suitable compounds of Formula I can be prepared by a method comprising the steps of:

- (a) treating with a strong base a lactone of the Formula A or B:




- 5       wherein R' and R<sup>2'</sup> are R and R<sup>2</sup>, respectively, or are suitably protected hydroxy groups; Ar<sup>10</sup> is Ar<sup>1</sup>, a suitably protected hydroxy-substituted aryl or a suitably protected amino-substituted aryl; and the remaining variables are as defined above for Formula I, provided that in lactone of formula B, when n and r are each zero, p is 1-4;

- 10       (b) reacting the product of step (a) with an imine of the formula




wherein Ar<sup>20</sup> is Ar<sup>2</sup>, a suitably protected hydroxy-substituted aryl or a suitably protected amino-substituted aryl; and Ar<sup>30</sup> is Ar<sup>3</sup>, a suitably protected hydroxy-substituted aryl or a suitably protected amino-substituted aryl;

- 15       c) quenching the reaction with an acid;
- d) optionally removing the protecting groups from R', R<sup>2'</sup>, Ar<sup>10</sup>, Ar<sup>20</sup> and Ar<sup>30</sup>, when present; and
- e) optionally functionalizing hydroxy or amino substituents at R, R<sup>2</sup>, Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup>.
- 20       Using the lactones shown above, compounds of Formula IA and IB are obtained as follows:




wherein the variables are as defined above; and



wherein the variables are as defined above.

- 5            Alternative sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (III) below:



10

(III)

- or isomers of the compounds of Formula (III), or pharmaceutically acceptable salts or solvates of the compounds of Formula (III) or of the isomers of the compounds of Formula (III), or prodrugs of the compounds of Formula (III) or of the isomers, salts or solvates of the compounds of Formula (III), wherein, in Formula (III) above:

- 15            Ar<sup>1</sup> is R<sup>3</sup>-substituted aryl;  
               Ar<sup>2</sup> is R<sup>4</sup>-substituted aryl;  
               Ar<sup>3</sup> is R<sup>5</sup>-substituted aryl;  
               Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-,  
 20            -CH(lower alkyl)- and -C(dilower alkyl)-;

A is selected from -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

R<sup>1</sup> is selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> and -O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl and aryl; or R<sup>1</sup> and R<sup>2</sup> together are =O;

5 q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;

R<sup>5</sup> is 1-3 substituents independently selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>9</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>-lower alkyl, -NR<sup>6</sup>SO<sub>2</sub>-aryl, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>-alkyl, S(O)<sub>0-2</sub>-aryl, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, o-halogeno, m-halogeno, o-lower alkyl, m-lower alkyl, -(lower alkylene)-COOR<sup>6</sup>, and -CH=CH-COOR<sup>6</sup>;

R<sup>3</sup> and R<sup>4</sup> are independently 1-3 substituents independently selected from the group consisting of R<sup>5</sup>, hydrogen, p-lower alkyl, aryl, -NO<sub>2</sub>, -CF<sub>3</sub> and p-halogeno;

R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and R<sup>9</sup> is lower alkyl, aryl or aryl-substituted lower alkyl.

Preferred compounds of Formula I include those in which Ar<sup>1</sup> is R<sup>3</sup>-substituted phenyl, especially (4-R<sup>3</sup>)-substituted phenyl. Ar<sup>2</sup> is preferably R<sup>4</sup>-substituted phenyl, especially (4-R<sup>4</sup>)-substituted phenyl. Ar<sup>3</sup> is preferably R<sup>5</sup>-substituted phenyl, especially (4-R<sup>5</sup>)-substituted phenyl. Mono-substitution of each of Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> is preferred.

Y and Z are each preferably -CH<sub>2</sub>-. R<sup>2</sup> is preferably hydrogen. R<sup>1</sup> is preferably -OR<sup>6</sup> wherein R<sup>6</sup> is hydrogen, or a group readily metabolizable to a hydroxyl (such as -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> and -O(CO)NR<sup>6</sup>R<sup>7</sup>, defined above). Also preferred are compounds wherein R<sup>1</sup> and R<sup>2</sup> together are =O.

The sum of q and p is preferably 1 or 2, more preferably 1. Preferred are compounds wherein p is zero and q is 1. More preferred are compounds wherein p is zero, q is 1, Y is -CH<sub>2</sub>- and R<sup>1</sup> is -OR<sup>6</sup>, especially when R<sup>6</sup> is hydrogen.

Another group of preferred compounds is that in which Ar<sup>1</sup> is R<sup>3</sup>-substituted phenyl, Ar<sup>2</sup> is R<sup>4</sup>-substituted phenyl and Ar<sup>3</sup> is R<sup>5</sup>-substituted phenyl.

Also preferred are compounds wherein Ar<sup>1</sup> is R<sup>3</sup>-substituted phenyl, Ar<sup>2</sup> is R<sup>4</sup>-substituted phenyl, Ar<sup>3</sup> is R<sup>5</sup>-substituted phenyl, and the sum of p and q is 1 or 2, especially 1. More preferred are compounds wherein Ar<sup>1</sup> is R<sup>3</sup>-substituted phenyl, Ar<sup>2</sup> is R<sup>4</sup>-substituted phenyl, Ar<sup>3</sup> is R<sup>5</sup>-substituted phenyl, p is zero and q is 1.

A is preferably -O-.


R<sup>3</sup> is preferably -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>-alkyl, S(O)<sub>0-2</sub>-aryl, NO<sub>2</sub> or halogeno. A more preferred definition for R<sup>3</sup> is halogeno, especially fluoro or chloro.

R<sup>4</sup> is preferably hydrogen, lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, COR<sup>6</sup> or halogeno, wherein R<sup>6</sup> and R<sup>7</sup> are preferably independently hydrogen or lower alkyl, and R<sup>9</sup> is preferably lower alkyl. A more preferred definition for R<sup>4</sup> is hydrogen or halogeno, especially fluoro or chloro.

R<sup>5</sup> is preferably -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, -(lower alkylene)-COOR<sup>6</sup> or -CH=CH-COOR<sup>6</sup>, wherein R<sup>6</sup> and R<sup>7</sup> are preferably independently hydrogen or lower alkyl, and R<sup>9</sup> is preferably lower alkyl. A more preferred definition for R<sup>5</sup> is -OR<sup>6</sup>, -(lower alkylene)-COOR<sup>6</sup> or -CH=CH-COOR<sup>6</sup>, wherein R<sup>6</sup> is preferably hydrogen or lower alkyl.

Methods for making compounds of Formula III are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,688,990, which is incorporated herein by reference.

In another embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (IV):



(IV)


or isomers of the compounds of Formula (IV), or pharmaceutically acceptable salts or solvates of the compounds of Formula (IV) or of the isomers of the compounds of Formula (IV), or prodrugs of the compounds of Formula (IV) or of the isomers, salts or solvates of the compounds of Formula (IV), wherein, in Formula (IV) above:

A is selected from the group consisting of  $R^2$ -substituted heterocycloalkyl,  $R^2$ -substituted heteroaryl,  $R^2$ -substituted benzofused heterocycloalkyl, and  $R^2$ -substituted benzofused heteroaryl;

$Ar^1$  is aryl or  $R^3$ -substituted aryl;

10  $Ar^2$  is aryl or  $R^4$ -substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the



$R^1$  is selected from the group consisting of:


15  $-(CH_2)_q-$ , wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

$-(CH_2)_e-G-(CH_2)_r-$ , wherein G is  $-O-$ ,  $-C(O)-$ , phenylene,  $-NR^8-$  or  $-S(O)_{0-2}-$ , e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

$-(C_2-C_6 \text{ alkenylene})-$ ; and

20  $-(CH_2)_f-V-(CH_2)_g-$ , wherein V is  $C_3-C_6$  cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

$R^5$  is selected from:




$R^6$  and  $R^7$  are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(C_1-C_6 \text{ alkyl})-$ ,  $-C(\text{di-}(C_1-C_6) \text{ alkyl})-$ ,  $-CH=CH-$  and

25  $-C(C_1-C_6 \text{ alkyl})=CH-$ ; or  $R^5$  together with an adjacent  $R^6$ , or  $R^5$  together with an adjacent  $R^7$ , form a  $-CH=CH-$  or a  $-CH=C(C_1-C_6 \text{ alkyl})-$  group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when  $R^6$  is  $-CH=CH-$  or  $-C(C_1-C_6 \text{ alkyl})=CH-$ , a is 1; provided that when  $R^7$  is -

CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>6</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>7</sup>'s can be the same or different;

and when Q is a bond, R<sup>1</sup> also can be selected from:



where M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)- and -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl);

R<sup>10</sup> and R<sup>12</sup> are independently selected from the group consisting of -OR<sup>14</sup>, -O(CO)R<sup>14</sup>, -O(CO)OR<sup>16</sup> and -O(CO)NR<sup>14</sup>R<sup>15</sup>;

R<sup>11</sup> and R<sup>13</sup> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>10</sup> and R<sup>11</sup> together are =O, or R<sup>12</sup> and R<sup>13</sup> together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;


v is 0 or 1;


j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

R<sup>2</sup> is 1-3 substituents on the ring carbon atoms selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>2</sub>-C<sub>10</sub>)alkenyl, (C<sub>2</sub>-C<sub>10</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkenyl, R<sup>17</sup>-substituted aryl, R<sup>17</sup>-substituted benzyl, R<sup>17</sup>-substituted benzyloxy, R<sup>17</sup>-substituted aryloxy, halogeno, -NR<sup>14</sup>R<sup>15</sup>, NR<sup>14</sup>R<sup>15</sup>(C<sub>1</sub>-C<sub>6</sub> alkylene)-, NR<sup>14</sup>R<sup>15</sup>C(O)(C<sub>1</sub>-C<sub>6</sub> alkylene)-, -NHC(O)R<sup>16</sup>, OH, C<sub>1</sub>-C<sub>6</sub> alkoxy, -OC(O)R<sup>16</sup>, -COR<sup>14</sup>, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, NO<sub>2</sub>, -S(O)<sub>0-2</sub>R<sup>16</sup>, -SO<sub>2</sub>NR<sup>14</sup>R<sup>15</sup> and



-(C<sub>1</sub>-C<sub>6</sub> alkylene)COOR<sup>14</sup>; when R<sup>2</sup> is a substituent on a heterocycloalkyl ring, R<sup>2</sup> is

as defined, or is =O or ; and, where R<sup>2</sup> is a substituent on a substitutable ring nitrogen, it is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, arylcarbonyl, hydroxy, -(CH<sub>2</sub>)<sub>1-6</sub>CONR<sup>18</sup>R<sup>18</sup>,



wherein J is -O-, -NH-, -NR<sup>18</sup>- or -CH<sub>2</sub>-;

R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>14</sup>, -O(CO)R<sup>14</sup>, -O(CO)OR<sup>16</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>14</sup>, -O(CO)NR<sup>14</sup>R<sup>15</sup>, -NR<sup>14</sup>R<sup>15</sup>,  
 10 -NR<sup>14</sup>(CO)R<sup>15</sup>, -NR<sup>14</sup>(CO)OR<sup>16</sup>, -NR<sup>14</sup>(CO)NR<sup>15</sup>R<sup>19</sup>, -NR<sup>14</sup>SO<sub>2</sub>R<sup>16</sup>, -COOR<sup>14</sup>, -CONR<sup>14</sup>R<sup>15</sup>, -COR<sup>14</sup>, -SO<sub>2</sub>NR<sup>14</sup>R<sup>15</sup>, S(O)<sub>0-2</sub>R<sup>16</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>14</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>14</sup>R<sup>15</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>14</sup>, -CH=CH-COOR<sup>14</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>8</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>14</sup> or -COOR<sup>14</sup>;

15 R<sup>9</sup> and R<sup>17</sup> are independently 1-3 groups independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>14</sup>R<sup>15</sup>, OH and halogeno;

R<sup>14</sup> and R<sup>15</sup> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

20 R<sup>16</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>17</sup>-substituted aryl;

R<sup>18</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

R<sup>19</sup> is hydrogen, hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

As used in Formula (IV) above, "A" is preferably an R<sup>2</sup>-substituted, 6-membered heterocycloalkyl ring containing 1 or 2 nitrogen atoms. Preferred  
 25 heterocycloalkyl rings are piperidiny, piperaziny and morpholiny groups. The ring

"A" is preferably joined to the phenyl ring through a ring nitrogen. Preferred R<sup>2</sup> substituents are hydrogen and lower alkyl. R<sup>19</sup> is preferably hydrogen.

Ar<sup>2</sup> is preferably phenyl or R<sup>4</sup>-phenyl, especially (4-R<sup>4</sup>)-substituted phenyl. Preferred definitions of R<sup>4</sup> are lower alkoxy, especially methoxy, and halogeno, especially fluoro.

Ar<sup>1</sup> is preferably phenyl or R<sup>3</sup>-substituted phenyl, especially (4-R<sup>3</sup>)-substituted phenyl.

There are several preferred definitions for the -R<sup>1</sup>-Q- combination of variables:

Q is a bond and R<sup>1</sup> is lower alkylene, preferably propylene;

Q is a spiro group as defined above, wherein preferably R<sup>6</sup> and R<sup>7</sup> are each ethylene and R<sup>5</sup> is  $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$  or  $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$ ;

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{10} \\ | \\ -\text{M}-\text{Y}_d-\text{C}-\text{Z}_h- \\ | \\ \text{R}^{11} \end{array}$  wherein the variables

are chosen such that R<sup>1</sup> is -O-CH<sub>2</sub>-CH(OH)-;

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{12} \qquad \text{R}^{10} \\ | \qquad \qquad | \\ -\text{X}_m-(\text{C})_s-\text{Y}_n-(\text{C})_t-\text{Z}_p- \\ | \qquad \qquad | \\ \text{R}^{13} \qquad \text{R}^{11} \end{array}$  wherein the


variables are chosen such that R<sup>1</sup> is -CH(OH)-(CH<sub>2</sub>)<sub>2</sub>-; and

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{10} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{11} \end{array}$  wherein the

variables are chosen such that R<sup>1</sup> is -CH(OH)-CH<sub>2</sub>-S(O)<sub>0-2</sub>-.

Methods for making compounds of Formula IV are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,656,624, which is incorporated herein by reference.

In another embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (V):



(V)

or isomers of the compounds of Formula (V), or pharmaceutically acceptable salts or  
 5 solvates of the compounds of Formula (V) or of the isomers of the compounds of  
 Formula (V), or prodrugs of the compounds of Formula (V) or of the isomers, salts or  
 solvates of the compounds of Formula (V), wherein, in Formula (V) above:

Ar<sup>1</sup> is aryl, R<sup>10</sup>-substituted aryl or heteroaryl;

Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

10 Ar<sup>3</sup> is aryl or R<sup>5</sup>-substituted aryl;

X and Y are independently selected from the group consisting of -CH<sub>2</sub>-,  
 -CH(lower alkyl)- and -C(dilower alkyl)-;

R is -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> or -O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>1</sup> is hydrogen, lower alkyl  
 or aryl; or R and R<sup>1</sup> together are =O;

15 q is 0 or 1;

r is 0, 1 or 2;

m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and  
 q is 1, 2, 3, 4 or 5;

R<sup>4</sup> is 1-5 substituents independently selected from the group consisting of  
 20 lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>,  
 -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>,  
 -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;


R<sup>5</sup> is 1-5 substituents independently selected from the group consisting of -  
 25 OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>,  
 -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -  
 COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>,  
 -CN, -NO<sub>2</sub>, halogen, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl; and

$R^{10}$  is 1-5 substituents independently selected from the group consisting of  
 5 lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  
 $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  
 $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $-S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-CF_3$ ,  $-CN$ ,  $-NO_2$  and halogen.

Within the scope of Formula V, there are included two preferred structures. In  
 10 Formula VA, q is zero and the remaining variables are as defined above, and in  
 Formula VB, q is 1 and the remaining variables are as defined above:



$R^4$ ,  $R^5$  and  $R^{10}$  are each preferably 1-3 independently selected substituents as  
 set forth above. Preferred are compounds of Formula (V) wherein  $Ar^1$  is phenyl,  $R^{10}$ -  
 15 substituted phenyl or thienyl, especially  $(4-R^{10})$ -substituted phenyl or thienyl.  $Ar^2$  is  
 preferably  $R^4$ -substituted phenyl, especially  $(4-R^4)$ -substituted phenyl.  $Ar^3$  is  
 preferably phenyl or  $R^5$ -substituted phenyl, especially  $(4-R^5)$ -substituted phenyl.  
 When  $Ar^1$  is  $R^{10}$ -substituted phenyl,  $R^{10}$  is preferably halogeno, especially fluoro.  
 When  $Ar^2$  is  $R^4$ -substituted phenyl,  $R^4$  is preferably  $-OR^6$ , especially wherein  $R^6$  is  
 20 hydrogen or lower alkyl. When  $Ar^3$  is  $R^5$ -substituted phenyl,  $R^5$  is preferably halogeno,  
 especially fluoro. Especially preferred are compounds of Formula (V) wherein  $Ar^1$  is  
 phenyl, 4-fluorophenyl or thienyl,  $Ar^2$  is 4-(alkoxy or hydroxy)phenyl, and  $Ar^3$  is phenyl  
 or 4-fluorophenyl.

X and Y are each preferably  $-CH_2-$ . The sum of m, n and q is preferably 2, 3 or  
 25 4, more preferably 2. When q is 1, n is preferably 1 to 5.


Preferences for X, Y,  $Ar^1$ ,  $Ar^2$  and  $Ar^3$  are the same in each of Formulae (VA)  
 and (VB).

In compounds of Formula (VA), the sum of m and n is preferably 2, 3 or 4, more preferably 2. Also preferred are compounds wherein the sum of m and n is 2, and r is 0 or 1.

In compounds of Formula (VB), the sum of m and n is preferably 1, 2 or 3, more preferably 1. Especially preferred are compounds wherein m is zero and n is 1.  $R^1$  is preferably hydrogen and R is preferably  $-OR^6$  wherein  $R^6$  is hydrogen, or a group readily metabolizable to a hydroxyl (such as  $-O(CO)R^6$ ,  $-O(CO)OR^9$  and  $-O(CO)NR^6R^7$ , defined above), or R and  $R^1$  together form a  $=O$  group.

Methods for making compounds of Formula V are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,624,920, which is incorporated herein by reference.

In another embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (VI):



(VI)

or isomers of the compounds of Formula (VI), or pharmaceutically acceptable salts or solvates of the compounds of Formula (VI) or of the isomers of the compounds of Formula (VI), or prodrugs of the compounds of Formula (VI) or of the isomers, salts or solvates of the compounds of Formula (VI), wherein in Formula (VI):  $R_1$  is

$-\overset{|}{\text{CH}}-$ ,  $-\overset{|}{\text{C}}(\text{lower alkyl})-$ ,  $-\overset{|}{\text{CF}}-$ ,  $-\overset{|}{\text{C}}(\text{OH})-$ ,  $-\overset{|}{\text{C}}(\text{C}_6\text{H}_5)-$ ,  $-\overset{|}{\text{C}}(\text{C}_6\text{H}_4-\text{R}_{15})-$ ,

$-\overset{|}{\text{N}}-$  or  $-\overset{|}{\text{N}}\text{O}^-$ ;

$R_2$  and  $R_3$  are independently selected from the group consisting of:  $-\text{CH}_2-$ ,  $-\text{CH}(\text{lower alkyl})-$ ,  $-\text{C}(\text{di-lower alkyl})-$ ,  $-\text{CH}=\text{CH}-$  and  $-\text{C}(\text{lower alkyl})=\text{CH}-$ ; or

R<sub>1</sub> together with an adjacent R<sub>2</sub>, or R<sub>1</sub> together with an adjacent R<sub>3</sub>, form a -CH=CH- or a -CH=C(lower alkyl)- group;

u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sub>2</sub> is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R<sub>3</sub> is -CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, the R<sub>2</sub>'s can be the same or different; and provided that when u is 2 or 3, the R<sub>3</sub>'s can be the same or different;

R<sub>4</sub> is selected from B-(CH<sub>2</sub>)<sub>m</sub>C(O)-, wherein m is 0, 1, 2, 3, 4 or 5;

B-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 0, 1, 2, 3, 4, 5 or 6;

10 B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein Z is -O-, -C(O)-, phenylene, -N(R<sub>8</sub>)- or -S(O)<sub>0-2</sub>-, e is 0, 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4, 5 or 6;

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-;

B-(C<sub>4</sub>-C<sub>6</sub> alkadienylene)-;

15 B-(CH<sub>2</sub>)<sub>t</sub>-Z-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1, 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0, 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6;

20 B-(CH<sub>2</sub>)<sub>t</sub>-V-(C<sub>2</sub>-C<sub>6</sub> alkenylene)- or


B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-V-(CH<sub>2</sub>)<sub>t</sub>-, wherein V and t are as defined above, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>a</sub>-Z-(CH<sub>2</sub>)<sub>b</sub>-V-(CH<sub>2</sub>)<sub>d</sub>-, wherein Z and V are as defined above and a, b and d are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3, 4, 5 or 6; or


25 T-(CH<sub>2</sub>)<sub>s</sub>-, wherein T is cycloalkyl of 3-6 carbon atoms and s is 0, 1, 2, 3, 4, 5 or 6; or

R<sub>1</sub> and R<sub>4</sub> together form the group  $\text{B}-\text{CH}=\overset{\text{I}}{\text{C}}-$  ;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of  
30 pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl, thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or



- W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy, alkoxycarbonylalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl
- 5 lower alkanedioyl, allyloxy, -CF<sub>3</sub>, -OCF<sub>3</sub>, benzyl, R<sub>7</sub>-benzyl, benzyloxy, R<sub>7</sub>-benzyloxy, phenoxy, R<sub>7</sub>-phenoxy, dioxolanyl, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, OH, halogeno, -CN, -N<sub>3</sub>, -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, R<sub>11</sub>O<sub>2</sub>SNH-, (R<sub>11</sub>O<sub>2</sub>S)<sub>2</sub>N-, -S(O)<sub>2</sub>NH<sub>2</sub>, -S(O)<sub>0-2</sub>R<sub>8</sub>, tert-butyldimethyl-silyloxymethyl, -C(O)R<sub>12</sub>,
- 10 -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -CH=CHC(O)R<sub>12</sub>, -lower alkylene-C(O)R<sub>12</sub>, R<sub>10</sub>C(O)(lower alkyleneoxy)-, N(R<sub>8</sub>)(R<sub>9</sub>)C(O)(lower alkyleneoxy)- and



for substitution on ring carbon atoms,

- and the substituents on the substituted heteroaryl ring nitrogen atoms, when present, are selected from the group consisting of lower alkyl, lower alkoxy,
- 15 -C(O)OR<sub>10</sub>, -C(O)R<sub>10</sub>, OH, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, -S(O)<sub>2</sub>NH<sub>2</sub> and 2-(trimethylsilyl)-ethoxymethyl;


R<sub>7</sub> is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -COOH, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), OH, and halogeno;

R<sub>8</sub> and R<sub>9</sub> are independently selected from H or lower alkyl;

- 20 R<sub>10</sub> is selected from lower alkyl, phenyl, R<sub>7</sub>-phenyl, benzyl or R<sub>7</sub>-benzyl;

R<sub>11</sub> is selected from OH, lower alkyl, phenyl, benzyl, R<sub>7</sub>-phenyl or R<sub>7</sub>-benzyl;

R<sub>12</sub> is selected from H, OH, alkoxy, phenoxy, benzyloxy,



, -N(R<sub>8</sub>)(R<sub>9</sub>), lower alkyl, phenyl or R<sub>7</sub>-phenyl;

R<sub>13</sub> is selected from -O-, -CH<sub>2</sub>-, -NH-, -N(lower alkyl)- or -NC(O)R<sub>19</sub>;

- R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of H
- 25 and the groups defined for W; or R<sub>15</sub> is hydrogen and R<sub>16</sub> and R<sub>17</sub>, together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;

R<sub>19</sub> is H, lower alkyl, phenyl or phenyl lower alkyl; and

R<sub>20</sub> and R<sub>21</sub> are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzofused heteroaryl, W-substituted benzofused heteroaryl and cyclopropyl, wherein heteroaryl  
 5 is as defined above.

One group of preferred compounds of Formula VI is that in which R<sub>21</sub> is selected from phenyl, W-substituted phenyl, indanyl, benzofuranyl, benzodioxolyl, tetrahydronaphthyl, pyridyl, pyrazinyl, pyrimidinyl, quinolyl or cyclopropyl, wherein W is lower alkyl, lower alkoxy, OH, halogeno, -N(R<sub>8</sub>)(R<sub>9</sub>),  
 10 -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, NO<sub>2</sub>, -CN, -N<sub>3</sub>, -SH, -S(O)<sub>0-2</sub>-(lower alkyl), -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -COR<sub>12</sub>, phenoxy, benzyloxy, -OCF<sub>3</sub>, -CH=C(O)R<sub>12</sub> or tert-butyl dimethylsilyloxy, wherein R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>12</sub> and R<sub>19</sub> are as defined for Formula IV. When W is 2 or 3 substituents, the substituents can be the same or different.

15 Another group of preferred compounds of Formula VI is that in which R<sub>20</sub> is phenyl or W-substituted phenyl, wherein preferred meanings of W are as defined above for preferred definitions of R<sub>21</sub>.


More preferred are compounds of Formula VI wherein R<sub>20</sub> is phenyl or W-substituted phenyl and R<sub>21</sub> is phenyl, W-substituted phenyl, indanyl, benzofuranyl, benzodioxolyl, tetrahydronaphthyl, pyridyl, pyrazinyl, pyrimidinyl, quinolyl or cyclopropyl; W is lower alkyl, lower alkoxy, OH, halogeno, -N(R<sub>8</sub>)(R<sub>9</sub>), -  
 20 NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, NO<sub>2</sub>, -CN, -N<sub>3</sub>, -SH, -S(O)<sub>0-2</sub>-(lower alkyl), -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -COR<sub>12</sub>, phenoxy, benzyloxy, -CH=CHC(O)R<sub>12</sub>, -OCF<sub>3</sub> or tert-butyl-dimethyl-silyloxy, wherein when W is 2 or 3  
 25 substituents, the substituents can be the same or different, and wherein R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>12</sub> and R<sub>19</sub> are as defined in Formula VI.

Also preferred are compounds of Formula VI wherein R<sub>1</sub> is  $\overset{|}{\text{CH-}}$  or  $\overset{|}{\text{C(OH)-}}$ .

Another group of preferred compounds of Formula VI is in which R<sub>2</sub> and R<sub>3</sub> are each -CH<sub>2</sub>- and the sum of u and v is 2, 3 or 4, with u=v=2 being more preferred.



R<sub>4</sub> is preferably B-(CH<sub>2</sub>)<sub>q</sub>- or B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein B, Z, q, e and r are



as defined above. B is preferably hydrogen and wherein R<sub>15</sub> is preferably H, OH, lower alkoxy, especially methoxy, or halogeno, especially chloro.

5 Preferably Z is -O-, e is 0, and r is 0.

Preferably q is 0-2.

R<sub>20</sub> is preferably phenyl or W-substituted phenyl.

Preferred W substituents for R<sub>20</sub> are lower alkoxy, especially methoxy and ethoxy, OH, and -C(O)R<sub>12</sub>, wherein R<sub>12</sub> is preferably lower alkoxy.


10 Preferably R<sub>21</sub> is selected from phenyl, lower alkoxy-substituted phenyl and F-phenyl.

Especially preferred are compounds of Formula VI wherein R<sub>1</sub> is -CH-, or

-C(OH)-, R<sub>2</sub> and R<sub>3</sub> are each -CH<sub>2</sub>-, u=v=2, R<sub>4</sub> is B-(CH<sub>2</sub>)<sub>q</sub>-, wherein B is phenyl or phenyl substituted by lower alkoxy or chloro, q is 0-2, R<sub>20</sub> is phenyl, OH-phenyl, lower  
15 alkoxy-substituted phenyl or lower alkoxycarbonyl-substituted phenyl, and R<sub>21</sub> is phenyl, lower alkoxy-substituted phenyl or F-phenyl.


Methods for making compounds of Formula VI are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,698,548, which is incorporated herein by reference.

20 In another embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formulae (VIIA) or (VIIB):



(VIIA)

25 or




(VIIB)

or isomers of the compounds of Formulae (VIIA) or (VIIB), or pharmaceutically acceptable salts or solvates of the compounds of Formulae (VIIA) or (VIIB) or of the isomers of the compounds of Formulae (VIIA) or (VIIB), or prodrugs of the compounds of Formulae (VIIA) or (VIIB) or of the isomers, salts or solvates of the compounds of Formulae (VIIA) or (VIIB), wherein in Formulae (VIIA) and (VIIB):


A is  $-\text{CH}=\text{CH}-$ ,  $-\text{C}\equiv\text{C}-$  or  $-(\text{CH}_2)_p-$  wherein p is 0, 1 or 2;

B is



10

B' is



D is  $-(\text{CH}_2)_m\text{C}(\text{O})-$  or  $-(\text{CH}_2)_q-$  wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;


E is  $\text{C}_{10}$  to  $\text{C}_{20}$  alkyl or  $-\text{C}(\text{O})-(\text{C}_9 \text{ to } \text{C}_{19})\text{-alkyl}$ , wherein the alkyl is straight or branched, saturated or containing one or more double bonds;

R is hydrogen,  $\text{C}_1\text{-C}_{15}$  alkyl, straight or branched, saturated or containing one or more double bonds, or  $\text{B}-(\text{CH}_2)_r-$ , wherein r is 0, 1, 2, or 3;

$\text{R}_1$ ,  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_1'$ ,  $\text{R}_2'$ , and  $\text{R}_3'$  are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, carboxy,  $\text{NO}_2$ ,  $\text{NH}_2$ , OH, halogeno, lower alkylamino, dilower alkylamino,  $-\text{NHC}(\text{O})\text{OR}_5$ ,  $\text{R}_6\text{O}_2\text{SNH}-$  and  $-\text{S}(\text{O})_2\text{NH}_2$ ;

20

$\text{R}_4$  is



wherein n is 0, 1, 2 or 3;

$R_5$  is lower alkyl; and

$R_6$  is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents  
 5 are 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino and dilower alkylamino.

Preferred are compounds of Formula (VIIA) wherein R is hydrogen, saturated or mono-unsaturated  $C_1$  - $C_{10}$  alkyl or phenyl. Another group of preferred compounds of Formula (VIIA) is that in which D is propyl (i.e.,  $-(CH_2)_q-$  and q is 3). A third group  
 10 of preferred compounds of Formula (VIIA) is that wherein  $R_4$  is p-methoxyphenyl or 2,4,6-trimethoxyphenyl. Still another group of preferred compounds of Formula (VIIA) is that wherein A is ethylene or a bond (i.e.,  $-(CH_2)_p-$  wherein p is zero).  $R_1'$ ,  $R_2'$ , and  $R_3'$  are preferably each hydrogen, and preferably  $R_1$  is hydrogen, hydroxy, nitro, lower alkoxy, amino or t-butoxycarbonyl-amino and  $R_2$  and  $R_3$  are each hydrogen.

15 More preferred are compounds of Formula (VIIA) wherein  $R_1'$ ,  $R_2'$ , and  $R_3'$  are each hydrogen;  $R_1$  is hydrogen, hydroxy, nitro, lower alkoxy, amino or t-butoxycarbonyl-amino and  $R_2$  and  $R_3$  are each hydrogen; R is hydrogen, ethyl or phenyl; D is propyl;  $R_4$  is p-methoxyphenyl or 2,4,6-trimethoxyphenyl; and A is ethylene or a bond.

20 Preferred compounds of Formula (VIIA), wherein  $B'$  is phenyl, are shown in the

following table:

D	R	A	B	R <sub>4</sub>
-(CH <sub>2</sub> ) <sub>3</sub> -	H	—	p-MeO-phenyl	p-MeO-phenyl
-CH <sub>2</sub> C(O)-	phenyl	—	phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	H	—	phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	H	—	p-OH-phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	H	ethylene	p-MeO-phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	H	—	3-MeO-phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	ethyl	—	phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	phenyl	—	phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	ethyl	—	phenyl	2,4,6-tri-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	methyl	—	phenyl	p-MeO-phenyl
-(CH <sub>2</sub> ) <sub>3</sub> -	H	—	p-NH <sub>2</sub> -phenyl	p-MeO-phenyl

The first-listed compound in the above table having the (3R,4S) absolute stereochemistry is more preferred.


Preferred compounds of Formula (VIIB) are those wherein R is hydrogen, methyl, ethyl, phenyl or phenylpropyl. Another group of preferred compounds of Formula (VIIB) is that wherein R<sub>4</sub> is p-methoxyphenyl or 2,4,6-trimethoxyphenyl. Still another group of preferred compounds of Formula (VIIB) is that wherein A is ethylene or a bond. Yet another group of preferred compounds of Formula (VIIB) is that wherein E is decyl, oleoyl or 7-Z-hexadecenyl. Preferably R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are each hydrogen.

More preferred compounds of Formula (VIIB) are those wherein R is hydrogen, methyl, ethyl, phenyl or phenylpropyl; R<sub>4</sub> is p-methoxyphenyl or 2,4,6-trimethoxyphenyl; A is ethylene or a bond; E is decyl, oleoyl or 7-Z-hexadecenyl; and R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are each hydrogen.

A preferred compound of Formula (VIIB) is that wherein E is decyl, R is hydrogen, B-A is phenyl and R<sub>4</sub> is p-methoxyphenyl.

In another embodiment, sterol absorption inhibitors useful in the compositions and methods of the present invention are represented by Formula (VIII):

5





(VIII)

or isomers of the compounds of Formula (VIII), or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIII) or of the isomers of the compounds of Formula (VIII), or prodrugs of the compounds of Formula (VIII) or of the isomers, salts or solvates of the compounds of Formula (VIII), wherein, in Formula (VIII) above,

R<sup>26</sup> is H or OG<sup>1</sup>;

G and G<sup>1</sup> are independently selected from the group consisting of



provided that when R<sup>26</sup> is H or

OH, G is not H;

15 R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy or -W-R<sup>30</sup>;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and

5 -C(O)aryl;

R<sup>30</sup> is selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

10 R<sup>31</sup> is selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;


T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyll or morpholinyll group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyll or morpholinyll group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

25 Q is a bond or, with the 3-position ring carbon of the azetidinone,

forms the spiro group ; and

R<sup>1</sup> is selected from the group consisting of

-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-E-(CH<sub>2</sub>)<sub>r</sub>-, wherein E is -O-, -C(O)-, phenylene, -NR<sup>22</sup>- or -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C<sub>2</sub>-C<sub>6</sub>)alkenylene-; and

-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R<sup>12</sup> is

-CH-, -C(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -CF-, -C(OH)-, -C(C<sub>6</sub>H<sub>4</sub>-R<sup>23</sup>)-, -N-, or -<sup>+</sup>N<sup>+</sup>O<sup>-</sup>;

R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl)-, -CH=CH- and  
 10 -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;


provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1;

provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1;

15 provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and

provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

and when Q is a bond, R<sup>1</sup> also can be:



M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

20 X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub>)alkyl- and -C(di-(C<sub>1</sub>-C<sub>6</sub>)alkyl);

R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>,  
 25 -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>,  
 -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>,  
 -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>,

-O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>,  
-CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>15</sup> and R<sup>17</sup> are independently selected from the group consisting of -OR<sup>19</sup>, -  
O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup> and -O(CO)NR<sup>19</sup>R<sup>20</sup>;

5 R<sup>16</sup> and R<sup>18</sup> are independently selected from the group consisting of H,  
(C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>15</sup> and R<sup>16</sup> together are =O, or R<sup>17</sup> and R<sup>18</sup> together are  
=O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

10 s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided  
that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

15 j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R<sup>1</sup> is 
$$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$$
, Ar<sup>1</sup> can also be  
pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl,  
pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-  
20 C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the  
group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>,

25 -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.



Ar<sup>2</sup> is preferably phenyl or R<sup>11</sup>-phenyl, especially (4-R<sup>11</sup>)-substituted phenyl. Preferred definitions of R<sup>11</sup> are lower alkoxy, especially methoxy, and halogeno, especially fluoro.

Ar<sup>1</sup> is preferably phenyl or R<sup>10</sup>-substituted phenyl, especially (4-R<sup>10</sup>)-substituted phenyl. Preferably R<sup>10</sup> is halogeno, and more preferably fluoro.

There are several preferred definitions for the -R<sup>1</sup>-Q- combination of variables:

Q is a bond and R<sup>1</sup> is lower alkylene, preferably propylene;

Q is a spiro group as defined above, wherein preferably R<sup>13</sup> and R<sup>14</sup> are each ethylene and R<sup>12</sup> is  $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$  or  $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$ , and R<sup>1</sup> is  $-(\text{CH}_2)_q$  wherein q is 0-6;

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{15} \\ | \\ -\text{M}-\text{Y}_d-\text{C}-\text{Z}_h- \\ | \\ \text{R}^{16} \end{array}$  wherein the variables

are chosen such that R<sup>1</sup> is -O-CH<sub>2</sub>-CH(OH)-;

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{17} \\ | \\ -\text{X}_m-(\text{C})_s-\text{Y}_n-(\text{C})_t-\text{Z}_p- \\ | \qquad \qquad | \\ \text{R}^{18} \qquad \qquad \text{R}^{16} \end{array}$  wherein the

variables are chosen such that R<sup>1</sup> is -CH(OH)-(CH<sub>2</sub>)<sub>2</sub>-; and

Q is a bond and R<sup>1</sup> is  $\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$  wherein the

variables are chosen such that R<sup>1</sup> is -CH(OH)-CH<sub>2</sub>-S(O)<sub>0-2</sub>-.

A preferred compound of Formula (VIII) therefore, is one wherein G and G<sup>1</sup> are as defined above and in which the remaining variables have the following definitions:

Ar<sup>1</sup> is phenyl or R<sup>10</sup>-substituted phenyl, wherein R<sup>10</sup> is halogeno;


Ar<sup>2</sup> is phenyl or R<sup>11</sup>-phenyl, wherein R<sup>11</sup> is 1 to 3 substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkoxy and halogeno;

Q is a bond and R<sup>1</sup> is lower alkylene; Q, with the 3-position

ring carbon of the azetidinone, forms the group  $(R^{12})_b(R^{13})_a$  wherein preferably  $R^{13}$  and  $R^{14}$  are each ethylene and  $a$  and  $b$  are each 1, and wherein  $R^{12}$  is

- $-\dot{\text{C}}\text{H}-$  or  $-\dot{\text{C}}(\text{OH})-$ ;  $Q$  is a bond and  $R^1$  is  $-\text{O}-\text{CH}_2-\text{CH}(\text{OH})-$ ;  $Q$  is a bond and  $R^1$  is  $-\text{CH}(\text{OH})-(\text{CH}_2)_2-$ ; or  $Q$  is a bond and  $R^1$  is  $-\text{CH}(\text{OH})-\text{CH}_2-\text{S}(\text{O})_0-2-$ .


Preferred variables for  $G$  and  $G^1$  groups of the formulae



are as follows:

- $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently selected from the group consisting of H, (C1-C6)alkyl, benzyl and acetyl.

Preferred variables for group  $G$  or  $G^1$  of the formula:



are as follows:

- $R^3$ ,  $R^{3a}$ ,  $R^4$  and  $R^{4a}$  are selected from the group consisting of H, (C1-C6)alkyl, benzyl and acetyl;

$R$ ,  $R^a$  and  $R^b$  are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C1-C6)alkoxy(C1-C6)alkoxy and -W- $R^{30}$ ,

- wherein  $W$  is  $-\text{O}-\text{C}(\text{O})-$  or  $-\text{O}-\text{C}(\text{O})-\text{NR}^{31}-$ ,  $R^{31}$  is H and  $R^{30}$  is (C1-C6)alkyl,  $-\text{C}(\text{O})-(\text{C1-C4})\text{alkoxy}-(\text{C1-C6})\text{alkyl}$ , T, T-(C1-C6)alkyl, or T or T-(C1-C6)alkyl wherein T is substituted by one or two halogeno or (C1-C6)alkyl groups.

Preferred R<sup>30</sup> substituents are selected from the group consisting of: 2-fluorophenyl, 2,4-difluoro-phenyl, 2,6-dichlorophenyl, 2-methylphenyl, 2-thienylmethyl, 2-methoxy-carbonylethyl, thiazol-2-yl-methyl, 2-furyl, 2-methoxycarbonylbutyl and phenyl.


5 Preferred combinations of R, R<sup>a</sup> and R<sup>b</sup> are as follows:

1) R, R<sup>a</sup> and R<sup>b</sup> are independently -OH or -O-C(O)-NH-R<sup>30</sup>, especially wherein R<sup>a</sup> is -OH and R and R<sup>b</sup> are -O-C(O)-NH-R<sup>30</sup> and R<sup>30</sup> is selected from the preferred substituents identified above, or wherein R and R<sup>a</sup> are each -OH and R<sup>b</sup> is -O-C(O)-NH-R<sup>30</sup> wherein R<sup>30</sup> is 2-fluorophenyl, 2,4-difluoro-phenyl, 2,6-dichlorophenyl;


2) R<sup>a</sup> is -OH, halogeno, azido or (C<sub>1</sub>-C<sub>6</sub>)-alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, R<sup>b</sup> is H, halogeno, azido or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and R is -O-C(O)-NH-R<sup>30</sup>, especially compounds wherein R<sup>a</sup> is -OH, R<sup>b</sup> is H and R<sup>30</sup> is 2-fluorophenyl;

3) R, R<sup>a</sup> and R<sup>b</sup> are independently -OH or -O-C(O)-R<sup>30</sup> and R<sup>30</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, T, or T substituted by one or two halogeno or (C<sub>1</sub>-C<sub>6</sub>)alkyl groups, especially compounds wherein R is -OH and R<sup>a</sup> and R<sup>b</sup> are -O-C(O)-R<sup>30</sup> wherein R<sup>30</sup> is 2-furyl; and

4) R, R<sup>a</sup> and R<sup>b</sup> are independently -OH or halogeno. Three additional classes of preferred compounds are those wherein the C<sup>1'</sup> anomeric oxy is beta, wherein the C<sup>2'</sup> anomeric oxy is beta, and wherein the R group is alpha. G and G<sup>1</sup> are preferably selected from:



25




5 wherein Ac is acetyl and Ph is phenyl.

Preferably, R<sup>26</sup> is H or OH, more preferably H. The -O-G substituent is preferably in the 4-position of the phenyl ring to which it is attached.

In another embodiment, sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are represented by

10 Formula (IX) below:




or isomers of the compounds of Formula (IX), or pharmaceutically acceptable salts or solvates of the compounds of Formula (IX) or of the isomers of the compounds of  
 15 Formula (IX), or prodrugs of the compounds of Formula (IX) or of the isomers, salts or solvates of the compounds of Formula (IX), wherein

R<sup>26</sup> is selected from the group consisting of:

- a) OH;
- b) OCH<sub>3</sub>;
- c) fluorine and
- d) chlorine.

20

R<sup>1</sup> is selected from the group consisting of



R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy and -W-R<sup>30</sup>;

W is independently selected from the group consisting of  
 5 -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

10 R<sup>30</sup> is independently selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is independently selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

15 T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;


R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of H, halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>,  
 20 (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent

bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;


5 Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, or, with the 3-position ring carbon of the azetidinone,



10 R<sup>12</sup> is



R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub> alkyl)), -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

15 a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1; provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

20 R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>, -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

5 R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

10 R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

Ar<sup>2</sup> is preferably phenyl or R<sup>11</sup>-phenyl, especially (4-R<sup>11</sup>)-substituted phenyl. Preferred definitions of R<sup>11</sup> are lower alkoxy, especially methoxy, and halogeno, especially fluoro.

15 Ar<sup>1</sup> is preferably phenyl or R<sup>10</sup>-substituted phenyl, especially (4-R<sup>10</sup>)-substituted phenyl. A preferred definition of R<sup>10</sup> is halogeno, especially fluoro.

Preferably Q is a lower alkyl or a spiro group as defined above, wherein

preferably R<sup>13</sup> and R<sup>14</sup> are each ethylene and R<sup>12</sup> is  $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$  or  $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$ .

20 A preferred compound of formula IX, therefore, is one wherein R<sup>1</sup> is as defined above and in which the remaining variables have the following definitions:

Ar<sup>1</sup> is phenyl or R<sup>10</sup>-substituted phenyl, wherein R<sup>10</sup> is halogeno;

Ar<sup>2</sup> is phenyl or R<sup>11</sup>-phenyl, wherein R<sup>11</sup> is 1 to 3 substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkoxy and halogeno;


Q is a lower alkyl (i.e. C-1 to C-2) with Q = C-2 being preferred, or Q, with the

25 3-position ring carbon of the azetidinone, forms the group  $\begin{array}{c} \diagup \\ \text{R}^{12} - \text{C} - \text{C} - \text{C} \\ | \quad | \quad | \\ \text{R}^{14})_b \quad \text{R}^{13})_a \end{array}$

wherein preferably R<sup>13</sup> and R<sup>14</sup> are each ethylene and a and b are each 1, and

wherein R<sup>12</sup> is  $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$  or  $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$ ;


Preferred variables for R<sup>1</sup> groups of the formula



are as follows:

$R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently selected from the group consisting of H, (C1-C6)alkyl, benzyl and acetyl.

5 Preferred variables for group  $R^1$  of the formula



are as follows:

$R^3$ ,  $R^{3a}$ ,  $R^4$  and  $R^{4a}$  are selected from the group consisting of H, (C1-C6)alkyl, benzyl and acetyl;

10  $R$ ,  $R^a$  and  $R^b$  are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C1-C6)alkoxy(C1-C6)alkoxy and -W- $R^{30}$ , wherein W is -O-C(O)- or -O-C(O)-NH- $R^{31}$ ,  $R^{31}$  is H and  $R^{30}$  is (C1-C6)alkyl, -C(O)-(C1-C4)alkoxy-(C1-C6)alkyl, T, T-(C1-C6)alkyl, or T or T-(C1-C6)alkyl wherein T is substituted by one or two halogeno or (C1-C6)alkyl groups.


15 Preferred  $R^{30}$  substituents are 2-fluorophenyl, 2,4-difluoro-phenyl, 2,6-dichlorophenyl, 2-methylphenyl, 2-thienylmethyl, 2-methoxy-carbonylethyl, thiazol-2-yl-methyl, 2-furyl, 2-methoxycarbonylbutyl and phenyl. Preferred combinations of  $R$ ,  $R^a$  and  $R^b$  are as follows: (1)  $R$ ,  $R^a$  and  $R^b$  are independently -OH or -O-C(O)-NH- $R^{30}$ , especially wherein  $R^a$  is -OH and  $R$  and  $R^b$  are -O-C(O)-NH- $R^{30}$  and  $R^{30}$  is selected from the preferred substituents identified above, or wherein  $R$  and  $R^a$  are -OH and  $R^b$  is -O-C(O)-NH- $R^{30}$  wherein  $R^{30}$  is 2-fluorophenyl, 2,4-difluoro-phenyl, 2,6-dichlorophenyl; (2)  $R^a$  is -OH, halogeno, azido or (C1-C6)-alkoxy(C1-C6)alkoxy,  $R^b$  is H, halogeno, azido or (C1-C6)alkoxy(C1-C6)-alkoxy, and  $R$  is -O-C(O)-NH- $R^{30}$ , especially compounds wherein  $R^a$  is -OH,  $R^b$  is H and  $R^{30}$  is 2-fluorophenyl; (3)  $R$ ,  $R^a$  and  $R^b$  are independently -OH or -O-C(O)- $R^{30}$  and  $R^{30}$  is (C1-C6)alkyl, T, or

25





- T substituted by one or two halogeno or (C<sub>1</sub>-C<sub>6</sub>)alkyl groups, especially compounds wherein R is -OH and R<sup>a</sup> and R<sup>b</sup> are -O-C(O)-R<sup>30</sup> wherein R<sup>30</sup> is 2-furyl; and (4) R, R<sup>a</sup> and R<sup>b</sup> are independently -OH or halogeno. Three additional classes of preferred compounds are those wherein the C<sup>1'</sup> anomeric oxy is beta, wherein the C<sup>2'</sup> anomeric oxy is beta, and wherein the R group is alpha.

R<sup>1</sup> is preferably selected from:




10



- 15 wherein Ac is acetyl and Ph is phenyl.


An example of a useful compound of this invention is one represented by the formula X:



X

or pharmaceutically acceptable salts or solvates of the compound of Formula (X), or prodrugs of the compound of Formula (X) or of the salts or solvates of the compound of Formula (X), wherein R<sup>1</sup> is defined as above.

A more preferred compound is one represented by formula XI:



(XI)

or pharmaceutically acceptable salts or solvates of the compound of Formula (XI), or prodrugs of the compound of Formula (XI) or of the salts or solvates of the compound of Formula (XI).

In another embodiment, compositions, pharmaceutical compositions, therapeutic combinations, kits and methods of treatment as described above are provided which comprise: (a) a first amount of at least one of nicotinic acid or derivatives thereof; and (b) a second amount of at least one substituted azetidinone compound or at least one substituted  $\beta$ -lactam compound, or isomers of the at least one substituted azetidinone compound or the at least one substituted  $\beta$ -lactam compound, or pharmaceutically acceptable salts or solvates of the at least one substituted azetidinone compound or the at least one substituted  $\beta$ -lactam compound or of the isomers of the at least one substituted azetidinone compound or the at least one substituted  $\beta$ -lactam compound, or prodrugs of the at least one substituted azetidinone compound or the at least one substituted  $\beta$ -lactam compound or of the isomers, salts or solvates of the at least one substituted azetidinone compound or the at least one substituted  $\beta$ -lactam compound, wherein the first amount and the second

amount together in their totality (whether administered concurrently or consecutively) comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal. Suitable substituted azetidinone compounds or substituted  $\beta$ -lactam compounds can be selected from any of the compounds discussed above in Formulae I-XI. Other useful substituted azetidinone compounds include N-sulfonyl-2-azetidinones such as are disclosed in U.S. Patent No. 4,983,597 and ethyl 4-(2-oxoazetidin-4-yl)phenoxy-alkanoates such as are disclosed in Ram et al., Indian J. Chem. Sect. B. 29B, 12 (1990), p. 1134-7, which are incorporated by reference herein.

The compounds of Formulae I-XI can be prepared by known methods, including the methods discussed above and, for example, WO 93/02048 describes the preparation of compounds wherein  $-R^1-Q-$  is alkylene, alkenylene or alkylene interrupted by a hetero atom, phenylene or cycloalkylene; WO 94/17038 describes the preparation of compounds wherein Q is a spirocyclic group; WO 95/08532 describes the preparation of compounds wherein  $-R^1-Q-$  is a hydroxy-substituted alkylene group; PCT/US95/03196 describes compounds wherein  $-R^1-Q-$  is a hydroxy-substituted alkylene attached to the  $Ar^1$  moiety through an  $-O-$  or  $S(O)_0-2-$  group; and U.S. Serial No. 08/463,619, filed June 5, 1995, describes the preparation of compounds wherein  $-R^1-Q-$  is a hydroxy-substituted alkylene group attached the azetidinone ring by a  $-S(O)_0-2-$  group.

The daily dose of the sterol absorption inhibitor(s) preferably ranges from about 0.1 to about 1000 mg per day, and more preferably about 0.25 to about 50 mg/day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

For administration of pharmaceutically acceptable salts of the above compounds, the weights indicated above refer to the weight of the acid equivalent or the base equivalent of the therapeutic compound derived from the salt.

In one embodiment, the compositions or therapeutic combinations can further comprise one or more pharmacological or therapeutic agents or drugs such as cholesterol biosynthesis inhibitors and/or lipid-lowering agents discussed below.

In another embodiment, the composition or treatment can further comprise one or more cholesterol biosynthesis inhibitors coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above.

5           Non-limiting examples of cholesterol biosynthesis inhibitors for use in the compositions, therapeutic combinations and methods of the present invention include competitive inhibitors of HMG CoA reductase, the rate-limiting step in cholesterol biosynthesis, squalene synthase inhibitors, squalene epoxidase inhibitors and mixtures thereof. Non-limiting examples of suitable HMG CoA reductase inhibitors  
10           include statins such as lovastatin (for example MEVACOR® which is available from Merck & Co.), pravastatin (for example PRAVACHOL® which is available from Bristol Meyers Squibb), fluvastatin, simvastatin (for example ZOCOR® which is available from Merck & Co.), atorvastatin, cerivastatin, CI-981, and pitavastatin (such as NK-104 of Negma Kowa of Japan); HMG CoA synthetase inhibitors, for example L-  
15           659,699 ((E,E)-11-[3'R-(hydroxy-methyl)-4'-oxo-2'R-oxetanyl]-3,5,7R-trimethyl-2,4-undecadienoic acid); squalene synthesis inhibitors, for example squalestatin 1; and squalene epoxidase inhibitors, for example, NB-598 ((E)-N-ethyl-N-(6,6-dimethyl-2-hepten-4-ynyl)-3-[(3,3'-bithiophen-5-yl)methoxy]benzene-methanamine hydrochloride) and other sterol biosynthesis inhibitors such as DMP-565. Preferred HMG CoA  
20           reductase inhibitors include lovastatin, pravastatin and simvastatin. The most preferred HMG CoA reductase inhibitor is simvastatin.

          Generally, a total daily dosage of cholesterol biosynthesis inhibitor(s) can range from about 0.1 to about 160 mg per day, and preferably about 0.2 to about 80 mg/day in single or 2-3 divided doses.

25           In another preferred embodiment, the composition or treatment comprises the compound of Formula (II) in combination with one or more of nicotinic acid or derivatives thereof and one or more cholesterol biosynthesis inhibitors. In this embodiment, preferably the nicotinic acid or derivatives thereof is selected from nicotinic acid or acipimox. Preferably the cholesterol biosynthesis inhibitor comprises  
30           one or more HMG CoA reductase inhibitors, such as, for example, lovastatin, pravastatin and/or simvastatin. More preferably, the composition or treatment comprises the compound of Formula (II) in combination with simvastatin and nicotinic acid or acipimox.

Also useful with the present invention are compositions or therapeutic combinations that can further comprise at least one (one or more) activators for peroxisome proliferator-activated receptors (PPAR). The activators act as agonists for the peroxisome proliferator-activated receptors. Three subtypes of PPAR have been identified, and these are designated as peroxisome proliferator-activated receptor alpha (PPAR $\alpha$ ), peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) and peroxisome proliferator-activated receptor delta (PPAR $\delta$ ). It should be noted that PPAR $\delta$  is also referred to in the literature as PPAR $\beta$  and as NUC1, and each of these names refers to the same receptor.

PPAR $\alpha$  regulates the metabolism of lipids. PPAR $\alpha$  is activated by fibrates and a number of medium and long-chain fatty acids, and it is involved in stimulating  $\beta$ -oxidation of fatty acids. The PPAR $\gamma$  receptor subtypes are involved in activating the program of adipocyte differentiation and are not involved in stimulating peroxisome proliferation in the liver. PPAR $\delta$  has been identified as being useful in increasing high density lipoprotein (HDL) levels in humans. See, e.g., WO 97/28149.

PPAR $\alpha$  activator compounds are useful for, among other things, lowering triglycerides, moderately lowering LDL levels and increasing HDL levels. Useful examples of PPAR $\alpha$  activators include fibrates.

Non-limiting examples of suitable fibric acid derivatives ("fibrates") include clofibrate (such as ethyl 2-(p-chlorophenoxy)-2-methyl-propionate, for example ATROMID-S® Capsules which are commercially available from Wyeth-Ayerst); gemfibrozil (such as 5-(2,5-dimethylphenoxy)-2,2-dimethylpentanoic acid, for example LOPID® tablets which are commercially available from Parke Davis); ciprofibrate (C.A.S. Registry No. 52214-84-3, see U.S. Patent No. 3,948,973 which is incorporated herein by reference); bezafibrate (C.A.S. Registry No. 41859-67-0, see U.S. Patent No. 3,781,328 which is incorporated herein by reference); clinofibrate (C.A.S. Registry No. 30299-08-2, see U.S. Patent No. 3,716,583 which is incorporated herein by reference); binifibrate (C.A.S. Registry No. 69047-39-8, see BE 884722 which is incorporated herein by reference); lifibrol (C.A.S. Registry No. 96609-16-4); fenofibrate (such as TRICOR® micronized fenofibrate (2-[4-(4-chlorobenzoyl)phenoxy]-2-methyl-propanoic acid, 1-methylethyl ester) which is commercially available from Abbott Laboratories or LIPANTHYL® micronized fenofibrate which is commercially available from Laboratoire Fournier, France) and mixtures thereof. These

compounds can be used in a variety of forms, including but not limited to acid form, salt form, racemates, enantiomers, zwitterions and tautomers.

Other examples of PPAR $\alpha$  activators useful with the practice of the present invention include suitable fluorophenyl compounds as disclosed in U.S. No. 6,028,109 which is incorporated herein by reference; certain substituted phenylpropionic compounds as disclosed in WO 00/75103 which is incorporated herein by reference; and PPAR $\alpha$  activator compounds as disclosed in WO 98/43081 which is incorporated herein by reference.

Non-limiting examples of suitable PPAR $\gamma$  activators include derivatives of glitazones or thiazolidinediones, such as, troglitazone (such as REZULIN $\text{\textcircled{R}}$  troglitazone (-5-[[4-[3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl] methyl]-2,4-thiazolidinedione) commercially available from Parke-Davis); rosiglitazone (such as AVANDIA $\text{\textcircled{R}}$  rosiglitazone maleate (-5-[[4-[2-(methyl-2-pyridinylamino)ethoxy] phenyl] methyl]-2,4-thiazolidinedione, (Z) -2-butenedioate) (1:1) commercially available from SmithKline Beecham) and pioglitazone (such as ACTOS $\text{\textsuperscript{TM}}$  pioglitazone hydrochloride (5-[[4-[2-(5-ethyl-2-pyridinyl)ethoxy]phenyl]methyl]-2,4-] thiazolidinedione monohydrochloride) commercially available from Takeda Pharmaceuticals). Other useful thiazolidinediones include ciglitazone, englitazone, darglitazone and BRL 49653 as disclosed in WO 98/05331 which is incorporated herein by reference; PPAR $\gamma$  activator compounds disclosed in WO 00/76488 which is incorporated herein by reference; and PPAR $\gamma$  activator compounds disclosed in U.S. Patent No. 5,994,554 which is incorporated herein by reference.

Other useful PPAR $\gamma$  activator compounds include certain acetylphenols as disclosed in U.S. Patent No. 5,859,051 which is incorporated herein by reference; certain quinoline phenyl compounds as disclosed in WO 99/20275 which is incorporated herein by reference; aryl compounds as disclosed by WO 99/38845 which is incorporated herein by reference; certain 1,4-disubstituted phenyl compounds as disclosed in WO 00/63161; certain aryl compounds as disclosed in WO 01/00579 which is incorporated herein by reference; benzoic acid compounds as disclosed in WO 01/12612 and WO 01/12187 which are incorporated herein by reference; and substituted 4-hydroxy-phenylallic acid compounds as disclosed in WO 97/31907 which is incorporated herein by reference.

PPAR $\delta$  compounds are useful for, among other things, lowering triglyceride levels or raising HDL levels. Non-limiting examples of PPAR $\delta$  activators include suitable thiazole and oxazole derivates, such as C.A.S. Registry No. 317318-32-4, as disclosed in WO 01/00603 which is incorporated herein by reference); certain fluoro, chloro or thio phenoxy phenylacetic acids as disclosed in WO 97/28149 which is incorporated herein by reference; suitable non- $\beta$ -oxidizable fatty acid analogues as disclosed in U.S. Patent No. 5,093,365 which is incorporated herein by reference; and PPAR $\delta$  compounds as disclosed in WO 99/04815 which is incorporated herein by reference.

Moreover, compounds that have multiple functionality for activating various combinations of PPAR $\alpha$ , PPAR $\gamma$  and PPAR $\delta$  are also useful with the practice of the present invention. Non-limiting examples include certain substituted aryl compounds as disclosed in U.S. Patent No. 6,248,781; WO 00/23416; WO 00/23415; WO 00/23425; WO 00/23445; WO 00/23451; and WO 00/63153, all of which are incorporated herein by reference, are described as being useful PPAR $\alpha$  and/or PPAR $\gamma$  activator compounds. Other non-limiting examples of useful PPAR $\alpha$  and/or PPAR $\gamma$  activator compounds include activator compounds as disclosed in WO 97/25042 which is incorporated herein by reference; activator compounds as disclosed in WO 00/63190 which is incorporated herein by reference; activator compounds as disclosed in WO 01/21181 which is incorporated herein by reference; biaryl-oxa(thia)zole compounds as disclosed in WO 01/16120 which is incorporated herein by reference; compounds as disclosed in WO 00/63196 and WO 00/63209 which are incorporated herein by reference; substituted 5-aryl-2,4-thiazolidinediones compounds as disclosed in U.S. Patent No. 6,008,237 which is incorporated herein by reference; arylthiazolidinedione and aryloxazolidinedione compounds as disclosed in WO 00/78312 and WO 00/78313G which are incorporated herein by reference; GW2331 or (2-(4-[difluorophenyl]-1heptylureido)ethyl]phenoxy)-2-methylbutyric compounds as disclosed in WO 98/05331 which is incorporated herein by reference; aryl compounds as disclosed in U.S. Patent No. 6,166,049 which is incorporated herein by reference; oxazole compounds as disclosed in WO 01/17994 which is incorporated herein by reference; and dithiolane compounds as disclosed in WO 01/25225 and WO 01/25226 which are incorporated herein by reference.

Other useful PPAR activator compounds include substituted benzylthiazolidine-2,4-dione compounds as disclosed in WO 01/14349, WO 01/14350 and WO/01/04351

which are incorporated herein by reference; mercaptocarboxylic compounds as disclosed in WO 00/50392 which is incorporated herein by reference; ascofuranone compounds as disclosed in WO 00/53563 which is incorporated herein by reference; carboxylic compounds as disclosed in WO 99/46232 which is incorporated herein by reference; compounds as disclosed in WO 99/12534 which is incorporated herein by reference; benzene compounds as disclosed in WO 99/15520 which is incorporated herein by reference; o-anisamide compounds as disclosed in WO 01/21578 which is incorporated herein by reference; and PPAR activator compounds as disclosed in WO 01/40192 which is incorporated herein by reference.

10       The peroxisome proliferator-activated receptor(s) activator(s) are administered in a therapeutically effective amount to treat the specified condition, for example in a daily dose preferably ranging from about 50 to about 3000 mg per day, and more preferably about 50 to about 2000 mg per day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is  
15       dependent on such factors as the potency of the compound administered, the age, weight, condition and response of the patient.

      In another alternative embodiment, the compositions, therapeutic combinations or methods of the present invention can further comprise one or more bile acid sequestrants (insoluble anion exchange resins), coadministered with or in combination  
20       with the nicotinic acid or derivative(s) thereof and sterol absorption inhibitor(s) discussed above.

      Bile acid sequestrants bind bile acids in the intestine, interrupting the enterohepatic circulation of bile acids and causing an increase in the faecal excretion of steroids. Use of bile acid sequestrants is desirable because of their non-systemic  
25       mode of action. Bile acid sequestrants can lower intrahepatic cholesterol and promote the synthesis of apo B/E (LDL) receptors which bind LDL from plasma to further reduce cholesterol levels in the blood.

      Non-limiting examples of suitable bile acid sequestrants include cholestyramine (a styrene-divinylbenzene copolymer containing quaternary ammonium cationic  
30       groups capable of binding bile acids, such as QUESTRAN® or QUESTRAN LIGHT® which are available from Bristol-Myers Squibb), colestipol (a copolymer of diethylenetriamine and 1-chloro-2,3-epoxypropane, such as COLESTID® tablets which are available from Pharmacia), colestevlam hydrochloride (such as WelChol® Tablets (poly(allylamine hydrochloride) cross-linked with epichlorohydrin and alkylated



with 1-bromodecane and (6-bromohexyl)-trimethylammonium bromide) which are available from Sankyo), water soluble derivatives such as 3,3-ioene, N-(cycloalkyl) alkylamines and poliglusam, insoluble quaternized polystyrenes, saponins and mixtures thereof. Other useful bile acid sequestrants are disclosed in PCT Patent  
5 Applications Nos. WO 97/11345 and WO 98/57652, and U.S. Patents Nos. 3,692,895 and 5,703,188 which are incorporated herein by reference. Suitable inorganic cholesterol sequestrants include bismuth salicylate plus montmorillonite clay, aluminum hydroxide and calcium carbonate antacids.

Generally, a total daily dosage of bile acid sequestrant(s) can range from about  
10 1 to about 50 grams per day, and preferably about 2 to about 16 grams per day in single or 2-4 divided doses.

In an alternative embodiment, the compositions or treatments of the present invention can further comprise one or more ileal bile acid transport ("IBAT") inhibitors (or apical sodium co-dependent bile acid transport ("ASBT") inhibitors) coadministered  
15 with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. The IBAT inhibitors can inhibit bile acid transport to reduce LDL cholesterol levels. Non-limiting examples of suitable IBAT inhibitors include benzothiepinines such as therapeutic compounds comprising a 2,3,4,5-tetrahydro-1-benzothiepine 1,1-dioxide structure such as are disclosed in PCT  
20 Patent Application WO 00/38727 which is incorporated herein by reference.

Generally, a total daily dosage of IBAT inhibitor(s) can range from about 0.01 to about 1000 mg/day, and preferably about 0.1 to about 50 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the  
25 present invention can further comprise one or more AcylCoA:Cholesterol O-acyltransferase ("ACAT") Inhibitors, which can reduce LDL and HDL levels, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. ACAT is an enzyme responsible for esterifying excess intracellular cholesterol and may reduce the synthesis of VLDL,  
30 which is a product of cholesterol esterification, and overproduction of apo B-100-containing lipoproteins.

Non-limiting examples of useful ACAT inhibitors include avasimibe ([2,4,6-tris(1-methylethyl)phenyl]acetyl)sulfamic acid, 2,6-bis(1-methylethyl)phenyl ester, formerly known as CI-1011), HL-004, lecimibide (DuP-128) and CL-277082 (N-(2,4-

difluorophenyl)-*N*-[[4-(2,2-dimethylpropyl)phenyl]methyl]-*N*-heptylurea). See P. Chang et al., "Current, New and Future Treatments in Dyslipidaemia and Atherosclerosis", Drugs 2000 Jul;60(1); 55-93, which is incorporated by reference herein.

Generally, a total daily dosage of ACAT inhibitor(s) can range from about 0.1 to about 1000 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise one or more Cholesteryl Ester Transfer Protein ("CETP") Inhibitors coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. CETP is responsible for the exchange or transfer of cholesteryl ester carrying HDL and triglycerides in VLDL.

Non-limiting examples of suitable CETP inhibitors are disclosed in PCT Patent Application No. WO 00/38721 and U.S. Patent No. 6,147,090, which are incorporated herein by reference. Pancreatic cholesteryl ester hydrolase (pCEH) inhibitors such as WAY-121898 also can be coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above.

Generally, a total daily dosage of CETP inhibitor(s) can range from about 0.01 to about 1000 mg/day, and preferably about 0.5 to about 20 mg/kg body weight/day in single or divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise probucol or derivatives thereof (such as AGI-1067 and other derivatives disclosed in U.S. Patents Nos. 6,121,319 and 6,147,250), which can reduce LDL and HDL levels, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above.

Generally, a total daily dosage of probucol or derivatives thereof can range from about 10 to about 2000 mg/day, and preferably about 500 to about 1500 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise low-density lipoprotein (LDL) receptor activators, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Non-limiting examples of suitable LDL-receptor activators include HOE-402, an imidazolidinyl-pyrimidine derivative that directly stimulates LDL receptor activity. See M. Huettinger et al.,

"Hypolipidemic activity of HOE-402 is Mediated by Stimulation of the LDL Receptor Pathway", Arterioscler. Thromb. 1993; 13:1005-12.

Generally, a total daily dosage of LDL receptor activator(s) can range from about 1 to about 1000 mg/day in single or 2-4 divided doses.

5 In another alternative embodiment, the compositions or treatments of the present invention can further comprise fish oil, which contains Omega 3 fatty acids (3-PUFA), which can reduce VLDL and triglyceride levels, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Generally, a total daily dosage of fish oil or Omega 3  
10 fatty acids can range from about 1 to about 30 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise natural water soluble fibers, such as psyllium, guar, oat and pectin, which can reduce cholesterol levels, coadministered with or in  
15 combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Generally, a total daily dosage of natural water soluble fibers can range from about 0.1 to about 10 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise plant sterols, plant stanols and/or fatty acid  
20 esters of plant stanols, such as sitostanol ester used in BENECOL® margarine, which can reduce cholesterol levels, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Generally, a total daily dosage of plant sterols, plant stanols and/or fatty acid esters of  
25 plant stanols can range from about 0.5 to about 20 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise antioxidants, such as probucol, tocopherol, ascorbic acid,  $\beta$ -carotene and selenium, or vitamins such as vitamin B<sub>6</sub> or vitamin B<sub>12</sub>,  
30 coadministered with or in combination with the of nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Generally, a total daily dosage of antioxidants or vitamins can range from about 0.05 to about 10 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise monocyte and macrophage inhibitors such as polyunsaturated fatty acids (PUFA), thyroid hormones including throxine analogues such as CGS-26214 (a thyroxine compound with a fluorinated ring), gene therapy and  
5 use of recombinant proteins such as recombinant apo E, coadministered with or in combination with the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) discussed above. Generally, a total daily dosage of these agents can range from about 0.01 to about 1000 mg/day in single or 2-4 divided doses.

Also useful with the present invention are compositions or therapeutic  
10 combinations which further comprise hormone replacement agents and compositions. Useful hormone agents and compositions for hormone replacement therapy of the present invention include androgens, estrogens, progestins, their pharmaceutically acceptable salts and derivatives thereof. Combinations of these agents and compositions are also useful.

15 The dosage of androgen and estrogen combinations vary, desirably from about 1 mg to about 4 mg androgen and from about 1 mg to about 3 mg estrogen. Examples include, but are not limited to, androgen and estrogen combinations such as the combination of esterified estrogens (sodium estrone sulfate and sodium equilin sulfate) and methyltestosterone (17-hydroxy-17-methyl-, (17B)- androst-4-en-3-one)  
20 available from Solvay Pharmaceuticals, Inc., Marietta, GA, under the tradename Estratest.

Estrogens and estrogen combinations may vary in dosage from about 0.01 mg up to 8 mg, desirably from about 0.3 mg to about 3.0 mg. Examples of useful estrogens and estrogen combinations include:

25 (a) the blend of nine (9) synthetic estrogenic substances including sodium estrone sulfate, sodium equilin sulfate, sodium 17  $\alpha$  -dihydroequilin sulfate, sodium 17  $\alpha$  -estradiol sulfate, sodium 17  $\beta$  -dihydroequilin sulfate, sodium 17  $\alpha$  -dihydroequilenin sulfate, sodium 17  $\beta$  -dihydroequilenin sulfate, sodium equilenin sulfate and sodium 17  $\beta$  -estradiol sulfate; available from Duramed Pharmaceuticals, Inc., Cincinnati, OH,  
30 under the tradename Cenestin;

(b) ethinyl estradiol (19-nor-17  $\alpha$  -pregna-1,3,5(10)-trien-20-yne-3,17-diol; available by Schering Plough Corporation, Kenilworth, NJ, under the tradename Estinyl;

(c) esterified estrogen combinations such as sodium estrone sulfate and sodium equilin sulfate; available from Solvay under the tradename Estratab and from Monarch Pharmaceuticals, Bristol, TN, under the tradename Menest;

(d) estropipate (piperazine estra-1,3,5(10)-trien-17-one, 3-(sulfooxy)-estrone sulfate); available from Pharmacia & Upjohn, Peapack, NJ, under the tradename Ogen and from Women First Health Care, Inc., San Diego, CA, under the tradename Ortho-Est; and

(e) conjugated estrogens (17  $\alpha$ -dihydroequilin, 17  $\alpha$ -estradiol, and 17  $\beta$ -dihydroequilin); available from Wyeth-Ayerst Pharmaceuticals, Philadelphia, PA, under the tradename Premarin.

Progestins and estrogens may also be administered with a variety of dosages, generally from about 0.05 to about 2.0 mg progestin and about 0.001 mg to about 2 mg estrogen, desirably from about 0.1 mg to about 1 mg progestin and about 0.01 mg to about 0.5 mg estrogen. Examples of progestin and estrogen combinations that may vary in dosage and regimen include:

(a) the combination of estradiol (estra-1, 3, 5 (10)-triene-3, 17  $\beta$ -diol hemihydrate) and norethindrone (17  $\beta$ -acetoxy-19-nor-17  $\alpha$ -pregn-4-en-20-yn-3-one); which is available from Pharmacia & Upjohn, Peapack, NJ, under the tradename Activella;

(b) the combination of levonorgestrel (d(-)-13  $\beta$ -ethyl-17  $\alpha$ -ethinyl-17  $\beta$ -hydroxygon-4-en-3-one) and ethinyl estradiol; available from Wyeth-Ayerst under the tradename Alesse, from Watson Laboratories, Inc., Corona, CA, under the tradenames Levora and Trivora, Monarch Pharmaceuticals, under the tradename Nordette, and from Wyeth-Ayerst under the tradename Triphasil;

(c) the combination of ethynodiol diacetate (19-nor-17  $\alpha$ -pregn-4-en-20-yne-3  $\beta$ , 17-diol diacetate) and ethinyl estradiol; available from G.D. Searle & Co., Chicago, IL, under the tradename Demulen and from Watson under the tradename Zovia;

(d) the combination of desogestrel (13-ethyl-11-methylene-18,19-dinor-17  $\alpha$ -pregn-4-en-20-yn-17-ol) and ethinyl estradiol; available from Organon under the tradenames Desogen and Mircette, and from Ortho-McNeil Pharmaceutical, Raritan, NJ, under the tradename Ortho-Cept;

(e) the combination of norethindrone and ethinyl estradiol; available from Parke-Davis, Morris Plains, NJ, under the tradenames Estrostep and femhrt, from

Watson under the tradenames Microgestin, Necon, and Tri-Norinyl, from Ortho-McNeil under the tradenames Modicon and Ortho-Novum, and from Warner Chilcott Laboratories, Rockaway, NJ, under the tradename Ovcon;

(f) the combination of norgestrel ( $(\pm)$ -13-ethyl-17-hydroxy-18, 19-dinor-17  $\alpha$ -preg-4-en-20-yn-3-one) and ethinyl estradiol; available from Wyeth-Ayerst under the tradenames Ovral and Lo/Ovral, and from Watson under the tradenames Ogestrel and Low-Ogestrel;

(g) the combination of norethindrone, ethinyl estradiol, and mestranol (3-methoxy-19-nor-17  $\alpha$ -pregna-1,3,5(10)-trien-20-yn-17-ol); available from Watson under the tradenames Brevicon and Norinyl;

(h) the combination of 17  $\beta$ -estradiol (estra-1,3,5(10)-triene-3,17  $\beta$ -diol) and micronized norgestimate (17  $\alpha$ -17-(Acetyloxy)-13-ethyl-18,19-dinorpregn-4-en-20-yn-3-one-3-oxime); available from Ortho-McNeil under the tradename Ortho-Prefest;

(i) the combination of norgestimate (18,19-dinor-17-pregn-4-en-20-yn-3-one, 17-(acetyloxy)-13-ethyl-,oxime, (17( $\alpha$ )-(+)-) and ethinyl estradiol; available from Ortho-McNeil under the tradenames Ortho Cyclen and Ortho Tri-Cyclen; and

(j) the combination of conjugated estrogens (sodium estrone sulfate and sodium equilin sulfate) and medroxyprogesterone acetate (20-dione, 17-(acetyloxy)-6-methyl-, (6( $\alpha$ ))- pregn-4-ene-3); available from Wyeth-Ayerst under the tradenames Premphase and Prempro.

In general, a dosage of progestins may vary from about .05 mg to about 10 mg or up to about 200 mg if microsize progesterone is administered. Examples of progestins include norethindrone; available from ESI Lederle, Inc., Philadelphia, PA, under the tradename Aygestin, from Ortho-McNeil under the tradename Micronor, and from Watson under the tradename Nor-QD; norgestrel; available from Wyeth-Ayerst under the tradename Ovrette; micronized progesterone (pregn-4-ene-3, 20-dione); available from Solvay under the tradename Prometrium; and medroxyprogesterone acetate; available from Pharmacia & Upjohn under the tradename Provera.

The compositions, therapeutic combinations or methods of the present invention can further comprise one or more obesity control medications. Useful obesity control medications include, but are not limited to, drugs that reduce energy intake or suppress appetite, drugs that increase energy expenditure and nutrient-partitioning agents. Suitable obesity control medications include, but are not limited to, noradrenergic agents (such as diethylpropion, mazindol, phenylpropanolamine,

phentermine, phendimetrazine, phendamine tartrate, methamphetamine, phendimetrazine and tartrate); serotonergic agents (such as sibutramine, fenfluramine, dexfenfluramine, fluoxetine, fluvoxamine and paroxetine); thermogenic agents (such as ephedrine, caffeine, theophylline, and selective  $\beta$ 3-adrenergic agonists); alpha-blocking agents; kainite or AMPA receptor antagonists; leptin-lipolysis stimulated receptors; phosphodiesterase enzyme inhibitors; compounds having nucleotide sequences of the mahogany gene; fibroblast growth factor-10 polypeptides; monoamine oxidase inhibitors (such as bexloxadone, moclobemide, brofaromine, phenoxathine, esuprone, befol, toloxatone, pirlindol, amiflamine, sercloremine, bazinaprine, lazabemide, milacemide and caroxazone); compounds for increasing lipid metabolism (such as evodiamine compounds); and lipase inhibitors (such as orlistat). Generally, a total dosage of the above-described obesity control medications can range from 1 to 3,000 mg/day, desirably from about 1 to 1,000 mg/day and more desirably from about 1 to 200 mg/day in single or 2-4 divided doses.

The compositions, therapeutic combinations or methods of the present invention can further comprise one or more blood modifiers which are chemically different from the substituted azetidinone and substituted  $\beta$ -lactam compounds (such as compounds I-XI above) and the bile acid sequestrants discussed above, for example, they contain one or more different atoms, have a different arrangement of atoms or a different number of one or more atoms than the sterol absorption inhibitor(s) or bile acid sequestrants discussed above. Useful blood modifiers include but are not limited to anti-coagulants (argatroban, bivalirudin, dalteparin sodium, desirudin, dicumarol, lyapolate sodium, nafamostat mesylate, phenprocoumon, tinzaparin sodium, warfarin sodium); antithrombotic (anagrelide hydrochloride, bivalirudin, cilostazol, dalteparin sodium, danaparoid sodium, dazoxiben hydrochloride, efegatran sulfate, enoxaparin sodium, fluretofen, ifetroban, ifetroban sodium, lamifiban, lotrafiban hydrochloride, napsagatran, orbofiban acetate, roxifiban acetate, sibrafiban, tinzaparin sodium, trifenagrel, abciximab, zolimomab aritox); fibrinogen receptor antagonists (roxifiban acetate, fradafiban, orbofiban, lotrafiban hydrochloride, tirofiban, xemilofiban, monoclonal antibody 7E3, sibrafiban); platelet inhibitors (cilostazol, clopidogrel bisulfate, epoprostenol, epoprostenol sodium, ticlopidine hydrochloride, aspirin, ibuprofen, naproxen, sulindae, idomethacin, mefenamate, droxicam, diclofenac, sulfipyrazone, piroxicam, dipyridamole); platelet aggregation inhibitors (acadesine, beraprost, beraprost sodium, ciprostone calcium,

itazigrel, lifarizine, lotrafiban hydrochloride, orbofiban acetate, oxagrelate, fradafiban, orbofiban, tirofiban, xemilofiban); hemorrhheologic agents (pentoxifylline); lipoprotein associated coagulation inhibitors; Factor VIIa inhibitors (4H-31-benzoxazin-4-ones, 4H-3,1-benzoxazin-4-thiones, quinazolin-4-ones, quinazolin-4-thiones, benzothiazin-4-ones, imidazolyl-boronic acid-derived peptide analogues TFPI-derived peptides, naphthalene-2-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolidin-3-(S)-yl} amide trifluoroacetate, dibenzofuran-2-sulfonic acid {1-[3-(aminomethyl)-benzyl]-5-oxo-pyrrolidin-3-yl}-amide, toluene-4-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolidin-3-(S)-yl}-amide trifluoroacetate, 3,4-dihydro-1H-isoquinoline-2-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolidin-3-(S)-yl}-amide trifluoroacetate); Factor Xa inhibitors (disubstituted pyrazolines, disubstituted triazolines, substituted n-[(aminoiminomethyl)phenyl] propylamides, substituted n-[(aminomethyl)phenyl] propylamides, tissue factor pathway inhibitor (TFPI), low molecular weight heparins, heparinoids, benzimidazolines, benzoxazolinones, benzopiperazinones, indanones, dibasic (amidinoaryl) propanoic acid derivatives, amidinophenyl-pyrrolidines, amidinophenyl-pyrrolines, amidinophenyl-isoxazolidines, amidinoindoles, amidinoazoles, bis-arylsulfonylaminobenzamide derivatives, peptidic Factor Xa inhibitors).

The compositions, therapeutic combinations or methods of the present invention can further comprise one or more cardiovascular agents which are chemically different from the substituted azetidinone and substituted  $\beta$ -lactam compounds (such as compounds I-XI above) and the bile acid sequestrants discussed above, for example, they contain one or more different atoms, have a different arrangement of atoms or a different number of one or more atoms than the sterol absorption inhibitor(s) or bile acid sequestrants discussed above. Useful cardiovascular agents include but are not limited to calcium channel blockers (clentiazem maleate, amlodipine besylate, isradipine, nimodipine, felodipine, nilvadipine, nifedipine, teludipine hydrochloride, diltiazem hydrochloride, belfosdil, verapamil hydrochloride, fostedil); adrenergic blockers (fenspiride hydrochloride, labetalol hydrochloride, proroxan, alfuzosin hydrochloride, acebutolol, acebutolol hydrochloride, alprenolol hydrochloride, atenolol, bunolol hydrochloride, carteolol hydrochloride, celiprolol hydrochloride, cetamolol hydrochloride, cicloprolol hydrochloride, dexpropranolol hydrochloride, diacetolol hydrochloride, dilevalol hydrochloride, esmolol hydrochloride, exaprolol hydrochloride, flestolol sulfate,



labetalol hydrochloride, levobetaxolol hydrochloride, levobunolol hydrochloride,  
 metalol hydrochloride, metoprolol, metoprolol tartrate, nadolol, pamatolol sulfate,  
 penbutolol sulfate, practolol, propranolol hydrochloride, sotalol hydrochloride, timolol,  
 timolol maleate, tiprenolol hydrochloride, tolamolol, bisoprolol, bisoprolol fumarate,  
 5   nebivolol); adrenergic stimulants; angiotensin converting enzyme (ACE) inhibitors  
 (benazepril hydrochloride, benazeprilat, captopril, delapril hydrochloride, fosinopril  
 sodium, libenzapril, moexipril hydrochloride, pentopril, perindopril, quinapril  
 hydrochloride, quinaprilat, ramipril, spirapril hydrochloride, spiraprilat, teprotide,  
 enalapril maleate, lisinopril, zofenopril calcium, perindopril erbumine); antihypertensive  
 10   agents (althiazide, benzthiazide, captopril, carvedilol, chlorothiazide sodium, clonidine  
 hydrochloride, cyclothiazide, delapril hydrochloride, dilevalol hydrochloride, doxazosin  
 mesylate, fosinopril sodium, guanfacine hydrochloride, methyldopa, metoprolol  
 succinate, moexipril hydrochloride, monatepil maleate, pelanserlin hydrochloride,  
 phenoxybenzamine hydrochloride, prazosin hydrochloride, primidolol, quinapril  
 15   hydrochloride, quinaprilat, ramipril, terazosin hydrochloride, candesartan, candesartan  
 cilexetil, telmisartan, amlodipine besylate, amlodipine maleate, bevantolol  
 hydrochloride); angiotensin II receptor antagonists (candesartan, irbesartan, losartan  
 potassium, candesartan cilexetil, telmisartan); anti-anginal agents (amlodipine  
 besylate, amlodipine maleate, betaxolol hydrochloride, bevantolol hydrochloride,  
 20   butopropazine hydrochloride, carvedilol, cinepazet maleate, metoprolol succinate,  
 molsidomine, monatepil maleate, primidolol, ranolazine hydrochloride, tofenfen,  
 verapamil hydrochloride); coronary vasodilators (fostedil, azaclozine hydrochloride,  
 chromonar hydrochloride, clonitrate, diltiazem hydrochloride, dipyridamole,  
 droprenilamine, erythryl tetranitrate, isosorbide dinitrate, isosorbide mononitrate,  
 25   lidoflazine, mioflazine hydrochloride, mixidine, molsidomine, nicorandil, nifedipine,  
 nisoldipine, nitroglycerine, oxprenolol hydrochloride, pentrinitrol, perhexiline maleate,  
 prenylamine, propatyl nitrate, terodiline hydrochloride, tolamolol, verapamil); diuretics  
 (the combination product of hydrochlorothiazide and spironolactone and the  
 combination product of hydrochlorothiazide and triamterene).

30       The compositions, therapeutic combinations or methods of the present  
 invention can further comprise one or more antidiabetic medications for reducing  
 blood glucose levels in a human. Useful antidiabetic medications include, but are not  
 limited to, drugs that reduce energy intake or suppress appetite, drugs that increase  
 energy expenditure and nutrient-partitioning agents. Suitable antidiabetic medications

include, but are not limited to, sulfonylurea (such as acetohexamide, chlorpropamide, gliamilide, gliclazide, glimepiride, glipizide, glyburide, glibenclamide, tolazamide, and tolbutamide), meglitinide (such as repaglinide and nateglinide), biguanide (such as metformin and buformin), alpha-glucosidase inhibitor (such as acarbose, miglitol, camiglibose, and voglibose), certain peptides (such as amlintide, pramlintide, exendin, and GLP-1 agonistic peptides), and orally administrable insulin or insulin composition for intestinal delivery thereof. Generally, a total dosage of the above-described antidiabetic medications can range from 0.1 to 1,000 mg/day in single or 2-4 divided doses.

10           Mixtures of any of the pharmacological or therapeutic agents described above can be used in the compositions and therapeutic combinations of the present invention.

          The compositions and therapeutic combinations of the present invention can be administered to a mammal in need of such treatment in a therapeutically effective amount to treat vascular conditions such as atherosclerosis, hyperlipidaemia (including but not limited to hypercholesterolaemia, hypertriglyceridaemia, sitosterolemia), stroke, diabetes, obesity, and/or reduce the level of sterol(s) in the plasma. The compositions and treatments can be administered by any suitable means which produce contact of these compounds with the site of action in the body, for example in the plasma, liver or ileum of a mammal or human.

20           The daily dosage for the various compositions and therapeutic combinations described above can be administered to a patient in a single dose or in multiple subdoses, as desired. Subdoses can be administered 2 to 6 times per day, for example. Sustained release dosages can be used. Where the nicotinic acid or derivatives thereof(s) and sterol absorption inhibitor(s) are administered in separate dosages, the number of doses of each component given per day may not necessarily be the same, e.g., one component may have a greater duration of activity and will therefore need to be administered less frequently.

          Since the present invention relates to reducing the plasma sterol (especially cholesterol) concentrations or levels by treatment with a combination of active ingredients wherein the active ingredients may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. That is, a kit is contemplated wherein two separate units are combined: a pharmaceutical composition comprising at least one of nicotinic acid or a derivative

thereof and a separate pharmaceutical composition comprising at least one sterol absorption inhibitor as described above. The kit will preferably include directions for the administration of the separate components. The kit form is particularly advantageous when the separate components must be administered in different  
5 dosage forms (e.g., oral and parenteral) or are administered at different dosage intervals.

The pharmaceutical treatment compositions and therapeutic combinations of the present invention can further comprise one or more pharmaceutically acceptable carriers, one or more excipients and/or one or more additives. Non-limiting examples  
10 of pharmaceutically acceptable carriers include solids and/or liquids such as ethanol, glycerol, water and the like. The amount of carrier in the treatment composition can range from about 5 to about 99 weight percent of the total weight of the treatment composition or therapeutic combination. Non-limiting examples of suitable pharmaceutically acceptable excipients and additives include non-toxic compatible  
15 fillers, binders such as starch, disintegrants, buffers, preservatives, anti-oxidants, lubricants, flavorings, thickeners, coloring agents, emulsifiers and the like. The amount of excipient or additive can range from about 0.1 to about 90 weight percent of the total weight of the treatment composition or therapeutic combination. One skilled in the art would understand that the amount of carrier(s), excipients and additives (if  
20 present) can vary.

The treatment compositions of the present invention can be administered in any conventional dosage form, preferably an oral dosage form such as a capsule, tablet, powder, cachet, suspension or solution. The formulations and pharmaceutical compositions can be prepared using conventional pharmaceutically acceptable and  
25 conventional techniques. Several examples of preparation of dosage formulations are provided below.

The following formulations exemplify some of the dosage forms of this invention. In each formulation, the term "Active Compound I" designates a substituted azetidinone compound,  $\beta$ -lactam compound or any of the compounds of Formulas I-  
30 VIII described herein above and the term "Active Compound II" designates nicotinic acid or derivatives thereof described herein above.

EXAMPLETablets

<u>No.</u>	<u>Ingredient</u>	<u>mg/tablet</u>
1	Active Compound I	10
2	Lactose monohydrate NF	55
3	Microcrystalline cellulose	20
4	Povidone (K29-32) USP	4
5	Croscarmellose sodium NF	8
6	Sodium lauryl sulfate	2
7	Magnesium stearate NF	1
	Total	100

5           In the present invention, the above-described tablet can be coadministered with a tablet, capsule, etc. comprising a dosage of Active Compound II, for example a NIASPAN® niacin extended-release tablet as described above.

Method of Manufacture

10           Mix Item No. 4 with purified water in suitable mixer to form binder solution. Spray the binder solution and then water over Items 1, 2, 6 and a portion of Item 5 in a fluidized bed processor to granulate the ingredients. Continue fluidization to dry the damp granules. Screen the dried granules and blend with Item No. 3 and the remainder of Item 5. Add Item No. 7 and mix. Compress the mixture to appropriate  
15 size and weight on a suitable tablet machine.

          For coadministration in separate tablets or capsules, representative formulations comprising a cholesterol absorption inhibitor such as are discussed above are well known in the art and representative formulations comprising a nicotinic acid derivative such as are discussed above are well known in the art. It is  
20 contemplated that where the two active ingredients are administered as a single composition, the dosage forms disclosed above for substituted azetidinone or  $\beta$ -lactam compounds may readily be modified using the knowledge of one skilled in the art.


The treatment compositions and therapeutic combinations of the present invention can inhibit the intestinal absorption of cholesterol in mammals, and can be useful in the treatment and/or prevention of vascular conditions, such as atherosclerosis, hypercholesterolemia and sitosterolemia, diabetes, obesity and  
5 lowering of plasma levels of cholesterol in mammals, in particular in mammals..

In another embodiment of the present invention, the compositions and therapeutic combinations of the present invention can inhibit sterol absorption or reduce plasma concentration of at least one sterol selected from the group consisting of phytosterols (such as sitosterol, campesterol, stigmasterol and avenosterol), 5 $\alpha$ -  
10 stanols (such as cholestanol, 5 $\alpha$ -campestanol, 5 $\alpha$ -sitostanol), cholesterol and mixtures thereof. The plasma concentration can be reduced by administering to a mammal in need of such treatment an effective amount of at least one treatment composition or therapeutic combination comprising at least one nicotinic acid or derivatives thereof and at least one sterol absorption inhibitor described above. The  
15 reduction in plasma concentration of sterols can range from about 1 to about 70 percent, and preferably about 10 to about 50 percent. Methods of measuring serum total blood cholesterol and total LDL cholesterol are well known to those skilled in the art and for example include those disclosed in PCT WO 99/38498 at page 11, incorporated by reference herein. Methods of determining levels of other sterols in  
20 serum are disclosed in H. Gylling et al., "Serum Sterols During Stanol Ester Feeding in a Mildly Hypercholesterolemic Population", J. Lipid Res. 40: 593-600 (1999), incorporated by reference herein.

Illustrating the invention are the following examples which; however, are not to be considered as limiting the invention to their details. Unless otherwise indicated, all  
25 parts and percentages in the following examples, as well as throughout the specification, are by weight.

### EXAMPLE 1

30 Hypercholesterolemic Golden Syrian hamsters were used to evaluate the *in vivo* efficacy of a cholesterol absorption inhibitor compound of Formula (XII):



(XII)

in combination with niacin. Compound XII can be prepared as shown in Example 9 of U.S. Patent No. 5,688,787, which is incorporated by reference herein.

5        Hamsters were fed a cholesterol-containing diet for 7 days, which resulted in a 2-fold increase in plasma cholesterol and a 20-fold increase in hepatic cholesteryl esters. It was hypothesized that a compound which blocks dietary cholesterol absorption would reduce the accumulation of hepatic cholesteryl esters and inhibit the increase in plasma total cholesterol levels, while niacin should reduce plasma  
10        triglyceride levels.

Male Golden Syrian hamsters (Charles River Labs, Wilmington, MA) weighing between 100 and 125g were fed Wayne rodent chow until study onset. At study onset (Day 1) animals were separated into groups (n=4-6/group) and fed chow supplemented with 0.5% by weight of cholesterol (Research Diets Inc., New  
15        Brunswick, NJ). The Control and Test compounds were administered once daily for 7 days, starting on Day 1 via oral gavage in 0.2 ml corn oil. Corn oil was used as the Control. The Test compounds included the compound of Formula (XII) (3mg/kg of body weight/day), niacin (100mg/kg of body weight/day), or the compound of Formula (XII) combined with niacin as described in Table 1 below.

20        On Day 7, blood was collected into tubes containing ethylenediaminetetraacetic acid (EDTA), and plasma was prepared by low speed centrifugation at 4°C. Liver samples (0.5g) were taken for neutral lipid analyses.

Nonfasted plasma cholesterol levels were determined by a modification of the cholesterol oxidase method, in which the reagents were available in a kit form from  
25        Wako Pure Chemicals Industries, Ltd. (Osaka, Japan). Ten µl of plasma was assayed for total cholesterol in 1 ml of 0.15 M tris buffer, pH 7.0 containing p-chlorophenol (0.1%), cholesterol oxidase (0.13 U/ml), cholesteryl ester hydrolase (0.13 U/ml), peroxidase (2.4 U/ml) and 4-aminoantipyrine (0.015%). Assays were carried out at 37°C for 10 min, along with cholesterol standards (Sigma Chemical Co., St. Louis,  
30        MO) and the resultant red quinone pigment's absorbance was determined spectrophotometrically at 505 nm.

Plasma triglyceride levels were determined enzymatically by a modification of the lipase-glycerol phosphate oxidase method, obtained in a kit form from Sigma Chemical Co. Ten  $\mu$ l of plasma and glycerol standards were added to 1 ml of buffer, pH 7.0 containing ATP (0.3 mmol/l), Mg (3.0 mmol), 4-aminoantipyrine (0.15 mmol/l),  
 5 Na N-ethyl-N-(3-sulfoethyl) n-anisidine (1.69 mmol/l), lipase (50,000 U/l), glycerol kinase (1000 U/l), glycerol phosphate oxidase (2000 U/l), and peroxidase (2000U/l), incubated at 37°C for 5 min., and the absorbance of the quinonemine dye was evaluated spectrophotometrically at 540 nm.

Samples of liver (0.5g) were lipid extracted. Lipid extracts were dried under nitrogen into HPLC sample vials, resuspended in hexane and injected onto a Zorbax  
 10 Sil (4.6 x 25 cm) silica column. Chromatography was performed using an isocratic mobile phase containing 98.8% hexane and 1.2% isopropanol at a flow rate of 2 ml/min. Lipids were detected by absorbance at 206 nm and quantitated by computer integration (System Gold, Beckman) of elution profiles. Elution time for cholesteryl  
 15 ester was 1.45 min. Cholesteryl ester content of liver-derived samples was derived from a standard curve constructed using known amounts of cholesteryl oleate. Cholesteryl oleate was used as the standard since this is the major cholesteryl ester species present in the liver and this specific cholesteryl ester has an extinction coefficient that approximates that of a weighted average for all the cholesteryl esters  
 20 present in the liver.

**Table 1**

Group	Plasma Cholesterol (mg/dl)	Plasma Triglyceride (mg/dl)	Hepatic Cholesteryl Ester (mg/g)
Control	283 $\pm$ 17	224 $\pm$ 21	24.94 $\pm$ 1.39
Compound XII 3mg/kg/day	214 $\pm$ 13	200 $\pm$ 30	12.75 $\pm$ 1.12
Niacin 100mg/kg/day	252 $\pm$ 17	169 $\pm$ 17	22.05 $\pm$ 1.39
Compound XII (3mg/kg/day) + Niacin (100mg/kg/day)	238 $\pm$ 6	164 $\pm$ 28	14.93 $\pm$ 2.04

25 As shown in Table 1, Compound XII reduced plasma cholesterol levels and the accumulation of hepatic cholesteryl esters in the cholesterol-fed hamsters. Niacin reduced plasma triglyceride levels, but did not significantly reduce the cholesterol levels. The combination of Compound XII and niacin resulted in reductions in plasma and hepatic cholesterol levels, as well as plasma triglycerides (Table 1). These  
 30 results indicate that the combination of the cholesterol absorption inhibitor of

Compound XII and niacin can have additive effects on treating hyperlipidemia in male Golden Syrian hamsters, by reducing both cholesterol and triglyceride levels.

## **EXAMPLE 2**

### **PREPARATION OF COMPOUND OF FORMULA (II)**

5

Step 1): To a solution of (S)-4-phenyl-2-oxazolidinone (41 g, 0.25 mol) in CH<sub>2</sub>Cl<sub>2</sub> (200 ml), was added 4-dimethylaminopyridine (2.5 g, 0.02 mol) and triethylamine (84.7 ml, 0.61 mol) and the reaction mixture was cooled to 0°C. Methyl-4-(chloroformyl)butyrate (50 g, 0.3 mol) was added as a solution in CH<sub>2</sub>Cl<sub>2</sub> (375 ml) dropwise over 1 h, and the reaction was allowed to warm to 22°C. After 17 h, water and H<sub>2</sub>SO<sub>4</sub> (2N, 100 ml), was added the layers were separated, and the organic layer was washed sequentially with NaOH (10%), NaCl (sat'd) and water. The organic layer was dried over MgSO<sub>4</sub> and concentrated to obtain a semicrystalline product.

10

Step 2): To a solution of TiCl<sub>4</sub> (18.2 ml, 0.165 mol) in CH<sub>2</sub>Cl<sub>2</sub> (600 ml) at 0°C, was added titanium isopropoxide (16.5 ml, 0.055 mol). After 15 min, the product of Step 1 (49.0 g, 0.17 mol) was added as a solution in CH<sub>2</sub>Cl<sub>2</sub> (100 ml). After 5 min., diisopropylethylamine (DIPEA) (65.2 ml, 0.37 mol) was added and the reaction mixture was stirred at 0°C for 1 h, the reaction mixture was cooled to -20°C, and 4-benzyloxybenzylidene(4-fluoro)aniline (114.3 g, 0.37 mol) was added as a solid. The reaction mixture was stirred vigorously for 4 h at -20°C, then acetic acid was added as a solution in CH<sub>2</sub>Cl<sub>2</sub> dropwise over 15 min, the reaction mixture was allowed to warm to 0°C, and H<sub>2</sub>SO<sub>4</sub> (2N) was added. The reaction mixture was stirred an additional 1 h, the layers were separated, washed with water, separated and the organic layer was dried. The crude product was crystallized from ethanol/water to obtain the pure intermediate.

15

20

25

Step 3): To a solution of the product of Step 2 (8.9 g, 14.9 mmol) in toluene (100 ml) at 50°C, was added N,O-bis(trimethylsilyl)acetamide (BSA) (7.50 ml, 30.3 mmol). After 0.5 h, solid TBAF (0.39 g, 1.5 mmol) was added and the reaction mixture stirred at 50°C for an additional 3 h. The reaction mixture was cooled to 22°C, CH<sub>3</sub>OH (10 ml), was added. The reaction mixture was washed with HCl (1N), NaHCO<sub>3</sub> (1N) and NaCl (sat'd.), and the organic layer was dried over MgSO<sub>4</sub>.

30



Step 4): To a solution of the product of Step 3 (0.94 g, 2.2 mmol) in CH<sub>3</sub>OH (3 ml), was added water (1 ml) and LiOH·H<sub>2</sub>O (102 mg, 2.4 mmole). The reaction mixture was stirred at 22°C for 1 h and then additional LiOH·H<sub>2</sub>O (54 mg, 1.3 mmole) was added. After a total of 2 h, HCl (1N) and EtOAc was added, the layers were separated, the organic layer was dried and concentrated in *vacuo*. To a solution of the resultant product (0.91 g, 2.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> at 22°C, was added ClCOCOCl (0.29 ml, 3.3 mmol) and the mixture stirred for 16 h. The solvent was removed in *vacuo*.

Step 5): To an efficiently stirred suspension of 4-fluorophenylzinc chloride (4.4 mmol) prepared from 4-fluorophenylmagnesium bromide (1M in THF, 4.4 ml, 4.4 mmol) and ZnCl<sub>2</sub> (0.6 g, 4.4 mmol) at 4°C, was added tetrakis(triphenylphosphine)palladium (0.25 g, 0.21 mmol) followed by the product of Step 4 (0.94 g, 2.2 mmol) as a solution in THF (2 ml). The reaction was stirred for 1 h at 0°C and then for 0.5 h at 22°C. HCl (1N, 5 ml) was added and the mixture was extracted with EtOAc. The organic layer was concentrated to an oil and purified by silica gel chromatography to obtain 1-(4-fluorophenyl)-4(S)-(4-hydroxyphenyl)-3(R)-(3-oxo-3-phenylpropyl)-2-azetidinone:

HRMS calc'd for C<sub>24</sub>H<sub>19</sub>F<sub>2</sub>NO<sub>3</sub> = 408.1429, found 408.1411.

Step 6): To the product of Step 5 (0.95 g, 1.91 mmol) in THF (3 ml), was added (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo-[1,2-c][1,3,2] oxazaborole (120 mg, 0.43 mmol) and the mixture was cooled to -20°C. After 5 min, borohydride-dimethylsulfide complex (2M in THF, 0.85 ml, 1.7 mmol) was added dropwise over 0.5 h. After a total of 1.5 h, CH<sub>3</sub>OH was added followed by HCl (1 N) and the reaction mixture was extracted with EtOAc to obtain 1-(4-fluorophenyl)-3(R)-[3(S)-(4-fluorophenyl)-3-hydroxypropyl]-4(S)-[4-(phenylmethoxy)phenyl]-2-azetidinone (compound 6A-1) as an oil. <sup>1</sup>H in CDCl<sub>3</sub> d H<sub>3</sub> = 4.68. J = 2.3 Hz. CI (M<sup>+</sup>H) 500.

Use of (S)-tetra-hydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo-[1,2-c][1,3,2] oxazaborole gives the corresponding 3(R)-hydroxypropyl azetidinone (compound 6B-1). <sup>1</sup>H in CDCl<sub>3</sub> d H<sub>3</sub> = 4.69. J = 2.3 Hz. CI (M<sup>+</sup>H) 500.

To a solution of compound 6A-1 (0.4 g, 0.8 mmol) in ethanol (2 ml), was added 10% Pd/C (0.03 g) and the reaction mixture was stirred under a pressure (60 psi) of H<sub>2</sub> gas for 16 h. The reaction mixture was filtered and the solvent was concentrated to

obtain compound 6A. Mp 164-166°C; Cl (M<sup>+</sup>H) 410.  $[\alpha]_D^{25} = -28.1^\circ$  (c 3, CH<sub>3</sub>OH)  
Elemental analysis calc'd for C<sub>24</sub>H<sub>21</sub>F<sub>2</sub>NO<sub>3</sub>: C 70.41; H 5.17; N 3.42; found C  
70.25; H 5.19; N 3.54.

Similarly treat compound 6B-1 to obtain compound 6B.

- 5 Mp 129.5-132.5°C; Cl (M<sup>+</sup>H) 410. Elemental analysis calc'd for C<sub>24</sub>H<sub>21</sub>F<sub>2</sub>NO<sub>3</sub>: C  
70.41; H 5.17; N 3.42; found C 70.30; H 5.14; N 3.52.

Step 6' (Alternative): To a solution of the product of Step 5 (0.14 g, 0.3 mmol)  
in ethanol (2 ml), was added 10% Pd/C (0.03 g) and the reaction was stirred under a  
pressure (60 psi) of H<sub>2</sub> gas for 16 h. The reaction mixture was filtered and the solvent  
10 was concentrated to afford a 1:1 mixture of compounds 6A and 6B.


It will be appreciated by those skilled in the art that changes could be made to  
the embodiments described above without departing from the broad inventive concept  
15 thereof. It is understood, therefore, that this invention is not limited to the particular  
embodiments disclosed, but it is intended to cover modifications which are within the  
spirit and scope of the invention, as defined by the appended claims.

THEREFORE, WE CLAIM:

1. A composition comprising:

(a) at least one of nicotinic acid or derivatives thereof; and

(b) at least one sterol absorption inhibitor represented by Formula (I):



(I)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (I) or of the isomers thereof, or prodrugs of the compounds of Formula (I) or of the isomers, salts or solvates thereof,

wherein in Formula (I) above:

$\text{Ar}^1$  and  $\text{Ar}^2$  are independently selected from the group consisting of aryl and  $\text{R}^4$ -substituted aryl;

$\text{Ar}^3$  is aryl or  $\text{R}^5$ -substituted aryl;

X, Y and Z are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{CH}(\text{lower alkyl})-$  and  $-\text{C}(\text{dilower alkyl})-$ ;

R and  $\text{R}^2$  are independently selected from the group consisting of  $-\text{OR}^6$ ,  $-\text{O}(\text{CO})\text{R}^6$ ,  $-\text{O}(\text{CO})\text{OR}^9$  and  $-\text{O}(\text{CO})\text{NR}^6\text{R}^7$ ;

$\text{R}^1$  and  $\text{R}^3$  are independently selected from the group consisting of hydrogen, lower alkyl and aryl;

q is 0 or 1;

r is 0 or 1;

m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

$R^4$  is 1-5 substituents independently selected from the group consisting of lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 5  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$ ,  $-CH=CH-COOR^6$ ,  $-CF_3$ ,  $-CN$ ,  $-NO_2$  and halogen;

$R^5$  is 1-5 substituents independently selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ ,  
 10  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.

15

2. The composition according to claim 1, wherein the at least one of nicotinic acid or derivatives thereof is selected from the group consisting of nicotinic acid, niceritrol, nicofuranose, acipimox and mixtures thereof.


20

3. The composition according to claim 2, wherein the at least one of nicotinic acid or derivatives thereof is nicotinic acid.

4. The composition according to claim 1, wherein the at least one of nicotinic acid or derivatives thereof is administered to a mammal in an amount ranging  
 25 from about 500 to about 10,000 milligrams of nicotinic acid or derivatives thereof per day.

5. The composition according to claim 1, wherein the sterol absorption inhibitor is represented by Formula (II) below:

30



or a pharmaceutically acceptable salt or solvate thereof, or prodrug of the compound of Formula (II) or of the salt or solvate thereof.

5           6.     The composition according to claim 1, wherein the at least one sterol absorption inhibitor is administered to a mammal in an amount ranging from about 0.1 to about 1000 milligrams of sterol absorption inhibitor per day.

7.     The composition according to claim 1, further comprising at least one  
10 cholesterol biosynthesis inhibitor.

8.     The composition according to claim 7, wherein the at least one cholesterol biosynthesis inhibitor comprises at least one HMG CoA reductase inhibitor.

15

9.     The composition according to claim 8, wherein the at least one HMG CoA reductase inhibitor is selected from the group consisting of lovastatin, pravastatin, fluvastatin, simvastatin, atorvastatin, cerivastatin and mixtures thereof.

20

10.    The composition according to claim 9, wherein the at least one HMG CoA reductase inhibitor is simvastatin.

11.    The composition according to claim 1, further comprising at least one PPAR receptor activator.

25

12.    The composition according to claim 11, wherein the PPAR receptor activator is at least one fibric acid derivative is selected from the group consisting of

fenofibrate, clofibrate, gemfibrozil, ciprofibrate, bezafibrate, clonofibrate, binifibrate, lifibrol and mixtures thereof.

13. The composition according to claim 12, wherein the at least one fibric  
5 acid derivative is fenofibrate.

14. The composition according to claim 1, further comprising at least one  
bile acid sequestrant.

10 15. The composition according to claim 14, wherein the at least one bile acid  
sequestrant is selected from the group consisting of cholestyramine and colestipol.

16. The composition according to claim 1, further comprising at least one  
15 AcylCoA:Cholesterol O-acyltransferase Inhibitor.

17. The composition according to claim 1, further comprising probucol or  
derivatives thereof.

18. The composition according to claim 1, further comprising at least one  
20 low-density lipoprotein receptor activator.

19. The composition according to claim 1, further comprising at least one  
Omega 3 fatty acid.

20 20. The composition according to claim 1, further comprising at least one  
natural water soluble fiber.

21. The composition according to claim 1, further comprising at least one of  
plant sterols, plant stanols or fatty acid esters of plant stanols.

22. The composition according to claim 1, further comprising at least one  
30 antioxidant or vitamin.

23. The composition according to claim 1, further comprising at least one hormone replacement therapy composition.

24. The composition according to claim 1, further comprising at least one  
5 obesity control medication.

25. The composition according to claim 1, further comprising at least one blood modifier different from the compound of Formula (I).

10 26. The composition according to claim 1, further comprising at least one cardiovascular agent different from the compound of Formula I.


27. The composition according to claim 1, further comprising at least one antidiabetic medication.

15 28. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 1 and a pharmaceutically acceptable carrier.

20 29. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 1.

25 30. The method according to claim 29, wherein the vascular condition is hyperlipidemia.


30 31. A composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) a compound represented by Formula (II) below:



(II)

or pharmaceutically acceptable salt or solvate thereof, or prodrug of the compound of Formula (II) or of the salt or solvate thereof.

- 5            32.    A therapeutic combination comprising: (a) a first amount of at least one of nicotinic acid or derivatives thereof; and (b) a second amount of a compound represented by Formula (II) below:



(II)


- 10           or pharmaceutically acceptable salt or solvate thereof, or prodrug of the compound of Formula (II) or of the salt or solvate thereof, wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

15

33.    A therapeutic combination comprising:

- (a)    a first amount of at least one of nicotinic acid or derivatives thereof; and
- (b)    a second amount of at least one sterol absorption inhibitor represented
- 20           by Formula (I):





(I)

- 5 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (I) or of the isomers thereof, or prodrugs of the compounds of Formula (I) or of the isomers, salts or solvates thereof, wherein in Formula (I) above:

Ar<sup>1</sup> and Ar<sup>2</sup> are independently selected from the group consisting of aryl and R<sup>4</sup>-substituted aryl;

- 10 Ar<sup>3</sup> is aryl or R<sup>5</sup>-substituted aryl;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(lower alkyl)- and -C(dilower alkyl)-;

R and R<sup>2</sup> are independently selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> and -O(CO)NR<sup>6</sup>R<sup>7</sup>;

- 15 R<sup>1</sup> and R<sup>3</sup> are independently selected from the group consisting of hydrogen, lower alkyl and aryl;

q is 0 or 1;

r is 0 or 1;

- 20 m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

- R<sup>4</sup> is 1-5 substituents independently selected from the group consisting of lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -(lower alkylene)COOR<sup>6</sup>, -CH=CH-COOR<sup>6</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;
- 25

$R^5$  is 1-5 substituents independently selected from the group consisting of -  
 $OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  $-NR^6R^7$ ,  
 $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  $-CONR^6R^7$ , -  
 $COR^6$ ,  $-SO_2NR^6R^7$ ,  $S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
5  $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-(\text{lower alkylene})COOR^6$  and  $-CH=CH-COOR^6$ ;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of  
hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl  
wherein the first amount and the second amount together comprise a therapeutically  
10 effective amount for the treatment or prevention of a vascular condition, diabetes,  
obesity or lowering a concentration of a sterol in plasma of a mammal.

34. A therapeutic combination according to claim 33, wherein the at least  
one of nicotinic acid or derivatives thereof is administered concomitantly with the at  
15 least one sterol absorption inhibitor.

35. A therapeutic combination according to claim 33, wherein the at least  
one of nicotinic acid or derivatives thereof and the at least one sterol absorption  
inhibitor are present in separate treatment compositions.

20

36. A method of treating or preventing a vascular condition, diabetes,  
obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the  
step of administering to a mammal in need of such treatment an effective amount of  
the therapeutic combination of claim 33.


25

37. A composition comprising:

(a) at least one of nicotinic acid or derivatives thereof; and

(b) at least one sterol absorption inhibitor represented by Formula (III):

30



(III)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 5 of Formula (III) or of the isomers thereof, or prodrugs of the compounds of Formula  
 (III) or of the isomers, salts or solvates thereof, wherein, in Formula (III) above:

Ar<sup>1</sup> is R<sup>3</sup>-substituted aryl;

Ar<sup>2</sup> is R<sup>4</sup>-substituted aryl;

Ar<sup>3</sup> is R<sup>5</sup>-substituted aryl;

10 Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-,  
 -CH(lower alkyl)- and -C(dilower alkyl)-;

A is selected from -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

R<sup>1</sup> is selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> and -  
 O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl and  
 15 aryl; or R<sup>1</sup> and R<sup>2</sup> together are =O;

q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;

R<sup>5</sup> is 1-3 substituents independently selected from the group consisting of -  
 OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>9</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>,  
 20 -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>-lower alkyl,  
 -NR<sup>6</sup>SO<sub>2</sub>-aryl, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>-alkyl, S(O)<sub>0-2</sub>-aryl,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, o-halogeno, m-halogeno,  
 o-lower alkyl, m-lower alkyl, -(lower alkylene)-COOR<sup>6</sup>, and -CH=CH-COOR<sup>6</sup>;

R<sup>3</sup> and R<sup>4</sup> are independently 1-3 substituents independently selected from the  
 25 group consisting of R<sup>5</sup>, hydrogen, p-lower alkyl, aryl, -NO<sub>2</sub>, -CF<sub>3</sub> and  
 p-halogeno;

R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of  
 hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and


$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl.

38. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 37 and a pharmaceutically acceptable carrier.

39. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 38.

40. A therapeutic combination comprising:

- (a) a first amount of at least one of nicotinic acid or derivatives thereof; and
- (b) a second amount of at least one sterol absorption inhibitor represented by Formula (III):



(III)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (III) or of the isomers thereof, or prodrugs of the compounds of Formula (III) or of the isomers, salts or solvates thereof, wherein, in Formula (III) above:

$\text{Ar}^1$  is  $R^3$ -substituted aryl;

$\text{Ar}^2$  is  $R^4$ -substituted aryl;

$\text{Ar}^3$  is  $R^5$ -substituted aryl;

Y and Z are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{CH}(\text{lower alkyl})-$  and  $-\text{C}(\text{dilower alkyl})-$ ;

A is selected from -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

R<sup>1</sup> is selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> and -O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl and aryl; or R<sup>1</sup> and R<sup>2</sup> together are =O;

5 q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;

R<sup>5</sup> is 1-3 substituents independently selected from the group consisting of -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>9</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>-lower alkyl, 10 -NR<sup>6</sup>SO<sub>2</sub>-aryl, -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>-alkyl, S(O)<sub>0-2</sub>-aryl, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, o-halogeno, m-halogeno, o-lower alkyl, m-lower alkyl, -(lower alkylene)-COOR<sup>6</sup>, and -CH=CH-COOR<sup>6</sup>;

R<sup>3</sup> and R<sup>4</sup> are independently 1-3 substituents independently selected from the group consisting of R<sup>5</sup>, hydrogen, p-lower alkyl, aryl, -NO<sub>2</sub>, -CF<sub>3</sub> and 15 p-halogeno;


R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

R<sup>9</sup> is lower alkyl, aryl or aryl-substituted lower alkyl, wherein the first amount and the second amount together comprise a therapeutically 20 effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

41. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the 25 step of administering to a mammal in need of such treatment an effective amount of the therapeutic combination of claim 40.

42. A composition comprising:

- (a) at least one of nicotinic acid or derivatives thereof; and
- 30 (b) at least one sterol absorption inhibitor represented by Formula (IV):



(IV)


- 5 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (IV) or of the isomers thereof, or prodrugs of the compounds of Formula (IV) or of the isomers, salts or solvates thereof, wherein, in Formula (IV) above:

A is selected from the group consisting of R<sup>2</sup>-substituted heterocycloalkyl, R<sup>2</sup>-substituted heteroaryl, R<sup>2</sup>-substituted benzofused heterocycloalkyl, and R<sup>2</sup>-substituted benzofused heteroaryl;

Ar<sup>1</sup> is aryl or R<sup>3</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the



- 15 R<sup>1</sup> is selected from the group consisting of:


-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-G-(CH<sub>2</sub>)<sub>r</sub>-, wherein G is -O-, -C(O)-, phenylene, -NR<sup>8</sup>- or -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

- 20 -(C<sub>2</sub>-C<sub>6</sub> alkenylene)-; and

-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R<sup>5</sup> is selected from:




- 25 R<sup>6</sup> and R<sup>7</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl)-, -CH=CH- and

-C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>5</sup> together with an adjacent R<sup>6</sup>, or R<sup>5</sup> together with an adjacent R<sup>7</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sup>6</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1; provided that when R<sup>7</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>6</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>7</sup>'s can be the same or different;

and when Q is a bond, R<sup>1</sup> also can be selected from:



where M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)- and -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl);

R<sup>10</sup> and R<sup>12</sup> are independently selected from the group consisting of -OR<sup>14</sup>, -O(CO)R<sup>14</sup>, -O(CO)OR<sup>16</sup> and -O(CO)NR<sup>14</sup>R<sup>15</sup>;

R<sup>11</sup> and R<sup>13</sup> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>10</sup> and R<sup>11</sup> together are =O, or R<sup>12</sup> and R<sup>13</sup> together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;


s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;


v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

R<sup>2</sup> is 1-3 substituents on the ring carbon atoms selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>2</sub>-C<sub>10</sub>)alkenyl, (C<sub>2</sub>-C<sub>10</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkenyl, R<sup>17</sup>-substituted aryl, R<sup>17</sup>-substituted benzyl, R<sup>17</sup>-substituted benzyloxy, R<sup>17</sup>-substituted aryloxy, halogeno, -NR<sup>14</sup>R<sup>15</sup>, NR<sup>14</sup>R<sup>15</sup>(C<sub>1</sub>-C<sub>6</sub> alkylene)-, NR<sup>14</sup>R<sup>15</sup>C(O)(C<sub>1</sub>-C<sub>6</sub> alkylene)-, -NHC(O)R<sup>16</sup>, OH,

$C_1-C_6$  alkoxy,  $-OC(O)R^{16}$ ,  $-COR^{14}$ , hydroxy( $C_1-C_6$ )alkyl,  
 ( $C_1-C_6$ )alkoxy( $C_1-C_6$ )alkyl,  $NO_2$ ,  $-S(O)_{0-2}R^{16}$ ,  $-SO_2NR^{14}R^{15}$  and  
 $-(C_1-C_6 \text{ alkylene})COOR^{14}$ ; when  $R^2$  is a substituent on a heterocycloalkyl ring,  $R^2$  is

as defined, or is  $=O$  or ; and, where  $R^2$  is a substituent on a  
 5 substitutable ring nitrogen, it is hydrogen, ( $C_1-C_6$ )alkyl, aryl, ( $C_1-C_6$ )alkoxy, aryloxy,  
 ( $C_1-C_6$ )alkylcarbonyl, arylcarbonyl, hydroxy,  $-(CH_2)_{1-6}CONR^{18}R^{18}$ ,



wherein J is  $-O-$ ,  $-NH-$ ,  $-NR^{18}-$  or  $-CH_2-$ ;

$R^3$  and  $R^4$  are independently selected from the group consisting of 1-3  
 10 substituents independently selected from the group consisting of ( $C_1-C_6$ )alkyl,  
 $-OR^{14}$ ,  $-O(CO)R^{14}$ ,  $-O(CO)OR^{16}$ ,  $-O(CH_2)_{1-5}OR^{14}$ ,  $-O(CO)NR^{14}R^{15}$ ,  $-NR^{14}R^{15}$ ,  
 $-NR^{14}(CO)R^{15}$ ,  $-NR^{14}(CO)OR^{16}$ ,  $-NR^{14}(CO)NR^{15}R^{19}$ ,  $-NR^{14}SO_2R^{16}$ ,  $-COOR^{14}$ ,  
 $-CONR^{14}R^{15}$ ,  $-COR^{14}$ ,  $-SO_2NR^{14}R^{15}$ ,  $S(O)_{0-2}R^{16}$ ,  $-O(CH_2)_{1-10}COOR^{14}$ ,  
 $-O(CH_2)_{1-10}CONR^{14}R^{15}$ ,  $-(C_1-C_6 \text{ alkylene})COOR^{14}$ ,  $-CH=CHCOOR^{14}$ ,  $-CF_3$ ,  $-CN$ ,  $-$   
 15  $NO_2$  and halogen;

$R^8$  is hydrogen, ( $C_1-C_6$ )alkyl, aryl ( $C_1-C_6$ )alkyl,  $-C(O)R^{14}$  or  $-COOR^{14}$ ;

$R^9$  and  $R^{17}$  are independently 1-3 groups independently selected from the  
 group consisting of hydrogen, ( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxy,  $-COOH$ ,  $NO_2$ ,  
 $-NR^{14}R^{15}$ ,  $OH$  and halogeno;

20  $R^{14}$  and  $R^{15}$  are independently selected from the group consisting of hydrogen,  
 ( $C_1-C_6$ )alkyl, aryl and aryl-substituted ( $C_1-C_6$ )alkyl;

$R^{16}$  is ( $C_1-C_6$ )alkyl, aryl or  $R^{17}$ -substituted aryl;

$R^{18}$  is hydrogen or ( $C_1-C_6$ )alkyl; and

$R^{19}$  is hydrogen, hydroxy or ( $C_1-C_6$ )alkoxy.

25



43. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 42 and a pharmaceutically acceptable carrier.

5

44. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 43.


10

45. A therapeutic combination comprising:

(a) a first amount of at least one of nicotinic acid or derivatives thereof; and

(b) a second amount of at least one sterol absorption inhibitor represented by Formula (IV):

15



(IV)

20 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (IV) or of the isomers thereof, or prodrugs of the compounds of Formula (IV) or of the isomers, salts or solvates thereof, wherein, in Formula (IV) above:


A is selected from the group consisting of R<sup>2</sup>-substituted heterocycloalkyl, R<sup>2</sup>-substituted heteroaryl, R<sup>2</sup>-substituted benzofused heterocycloalkyl, and R<sup>2</sup>-substituted benzofused heteroaryl;

25

Ar<sup>1</sup> is aryl or R<sup>3</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the



$R^1$  is selected from the group consisting of:


$-(CH_2)_q-$ , wherein  $q$  is 2-6, provided that when  $Q$  forms a spiro ring,  $q$  can also be zero or 1;

$-(CH_2)_e-G-(CH_2)_r-$ , wherein  $G$  is  $-O-$ ,  $-C(O)-$ , phenylene,  $-NR^8-$  or  $-S(O)_{0-2}-$ ,  $e$  is 0-5 and  $r$  is 0-5, provided that the sum of  $e$  and  $r$  is 1-6;

$-(C_2-C_6 \text{ alkenylene})-$ ; and

$-(CH_2)_f-V-(CH_2)_g-$ , wherein  $V$  is  $C_3-C_6$  cycloalkylene,  $f$  is 1-5 and  $g$  is 0-5, provided that the sum of  $f$  and  $g$  is 1-6;


$R^5$  is selected from:



$R^6$  and  $R^7$  are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(C_1-C_6 \text{ alkyl})-$ ,  $-C(\text{di-}(C_1-C_6) \text{ alkyl})-$ ,  $-CH=CH-$  and  $-C(C_1-C_6 \text{ alkyl})=CH-$ ; or  $R^5$  together with an adjacent  $R^6$ , or  $R^5$  together with an adjacent  $R^7$ , form a  $-CH=CH-$  or a  $-CH=C(C_1-C_6 \text{ alkyl})-$  group;

$a$  and  $b$  are independently 0, 1, 2 or 3, provided both are not zero; provided that when  $R^6$  is  $-CH=CH-$  or  $-C(C_1-C_6 \text{ alkyl})=CH-$ ,  $a$  is 1; provided that when  $R^7$  is  $-CH=CH-$  or  $-C(C_1-C_6 \text{ alkyl})=CH-$ ,  $b$  is 1; provided that when  $a$  is 2 or 3, the  $R^6$ 's can be the same or different; and provided that when  $b$  is 2 or 3, the  $R^7$ 's can be the same or different;

and when  $Q$  is a bond,  $R^1$  also can be selected from:



where  $M$  is  $-O-$ ,  $-S-$ ,  $-S(O)-$  or  $-S(O)_2-$ ;

$X$ ,  $Y$  and  $Z$  are independently selected from the group consisting of  $-CH_2-$ ,  $-CH(C_1-C_6 \text{ alkyl})-$  and  $-C(\text{di-}(C_1-C_6) \text{ alkyl})-$ ;

$R^{10}$  and  $R^{12}$  are independently selected from the group consisting of  
 $-OR^{14}$ ,  $-O(CO)R^{14}$ ,  $-O(CO)OR^{16}$  and  $-O(CO)NR^{14}R^{15}$ ;

$R^{11}$  and  $R^{13}$  are independently selected from the group consisting of hydrogen,  
 $(C_1-C_6)$ alkyl and aryl; or  $R^{10}$  and  $R^{11}$  together are  $=O$ , or  $R^{12}$  and  $R^{13}$  together are  $=O$ ;

5 d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least  
 one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0  
 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the  
 10 sum of m, t and n is 1-5;


v is 0 or 1;


j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

$R^2$  is 1-3 substituents on the ring carbon atoms selected from the group  
 consisting of hydrogen,  $(C_1-C_{10})$ alkyl,  $(C_2-C_{10})$ alkenyl,  $(C_2-C_{10})$ alkynyl,

15  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkenyl,  $R^{17}$ -substituted aryl,  $R^{17}$ -substituted benzyl,  
 $R^{17}$ -substituted benzyloxy,  $R^{17}$ -substituted aryloxy, halogeno,  $-NR^{14}R^{15}$ ,  $NR^{14}R^{15}(C_1-$   
 $C_6$  alkylene)-,  $NR^{14}R^{15}C(O)(C_1-C_6$  alkylene)-,  $-NHC(O)R^{16}$ , OH,  
 $C_1-C_6$  alkoxy,  $-OC(O)R^{16}$ ,  $-COR^{14}$ , hydroxy $(C_1-C_6)$ alkyl,  
 $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,  $NO_2$ ,  $-S(O)_{0-2}R^{16}$ ,  $-SO_2NR^{14}R^{15}$  and

20  $-(C_1-C_6$  alkylene) $COOR^{14}$ ; when  $R^2$  is a substituent on a heterocycloalkyl ring,  $R^2$  is

as defined, or is  $=O$  or ; and, where  $R^2$  is a substituent on a  
 substitutable ring nitrogen, it is hydrogen,  $(C_1-C_6)$ alkyl, aryl,  $(C_1-C_6)$ alkoxy, aryloxy,  
 $(C_1-C_6)$ alkylcarbonyl, arylcarbonyl, hydroxy,  $-(CH_2)_{1-6}CONR^{18}R^{18}$ ,



25 wherein J is  $-O-$ ,  $-NH-$ ,  $-NR^{18}-$  or  $-CH_2-$ ;

$R^3$  and  $R^4$  are independently selected from the group consisting of 1-3  
 substituents independently selected from the group consisting of  $(C_1-C_6)$ alkyl,

-OR<sup>14</sup>, -O(CO)R<sup>14</sup>, -O(CO)OR<sup>16</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>14</sup>, -O(CO)NR<sup>14</sup>R<sup>15</sup>, -NR<sup>14</sup>R<sup>15</sup>,  
 -NR<sup>14</sup>(CO)R<sup>15</sup>, -NR<sup>14</sup>(CO)OR<sup>16</sup>, -NR<sup>14</sup>(CO)NR<sup>15</sup>R<sup>19</sup>, -NR<sup>14</sup>SO<sub>2</sub>R<sup>16</sup>, -COOR<sup>14</sup>,  
 -CONR<sup>14</sup>R<sup>15</sup>, -COR<sup>14</sup>, -SO<sub>2</sub>NR<sup>14</sup>R<sup>15</sup>, S(O)<sub>0-2</sub>R<sup>16</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>14</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>14</sup>R<sup>15</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>14</sup>, -CH=CH-COOR<sup>14</sup>, -CF<sub>3</sub>, -CN, -

5 NO<sub>2</sub> and halogen;

R<sup>8</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>14</sup> or -COOR<sup>14</sup>;

R<sup>9</sup> and R<sup>17</sup> are independently 1-3 groups independently selected from the  
 group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>,  
 -NR<sup>14</sup>R<sup>15</sup>, OH and halogeno;

10 R<sup>14</sup> and R<sup>15</sup> are independently selected from the group consisting of hydrogen,  
 (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>16</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>17</sup>-substituted aryl;

R<sup>18</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

R<sup>19</sup> is hydrogen, hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy,


15 wherein the first amount and the second amount together comprise a therapeutically  
 effective amount for the treatment or prevention of a vascular condition, diabetes,  
 obesity or lowering a concentration of a sterol in plasma of a mammal.

46. A method of treating or preventing a vascular condition, diabetes,  
 20 obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the  
 step of administering to a mammal in need of such treatment an effective amount of  
 the therapeutic combination of claim 45.

47. A composition comprising:

25 (a) at least one of nicotinic acid or derivatives thereof; and

(b) at least one sterol absorption inhibitor represented by Formula (V):



(V)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 5 of Formula (V) or of the isomers thereof, or prodrugs of the compounds of Formula (V)  
 or of the isomers, salts or solvates thereof, wherein, in Formula (V) above:

Ar<sup>1</sup> is aryl, R<sup>10</sup>-substituted aryl or heteroaryl;

Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

Ar<sup>3</sup> is aryl or R<sup>5</sup>-substituted aryl;

10 X and Y are independently selected from the group consisting of -CH<sub>2</sub>-,  
 -CH(lower alkyl)- and -C(dilower alkyl)-;

R is -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> or -O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>1</sup> is hydrogen, lower alkyl  
 or aryl; or R and R<sup>1</sup> together are =O;

q is 0 or 1;

15 r is 0, 1 or 2;

m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and  
 q is 1, 2, 3, 4 or 5;

R<sup>4</sup> is 1-5 substituents independently selected from the group consisting of  
 lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>,  
 20 -NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>,  
 -CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>,  
 -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

R<sup>5</sup> is 1-5 substituents independently selected from the group consisting of -  
 OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>,  
 25 -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -  
 COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -  
 NO<sub>2</sub>, halogen, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

$R^6$ ,  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;

$R^9$  is lower alkyl, aryl or aryl-substituted lower alkyl; and


$R^{10}$  is 1-5 substituents independently selected from the group consisting of  
 5 lower alkyl,  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$ ,  $-O(CH_2)_{1-5}OR^6$ ,  $-O(CO)NR^6R^7$ ,  
 $-NR^6R^7$ ,  $-NR^6(CO)R^7$ ,  $-NR^6(CO)OR^9$ ,  $-NR^6(CO)NR^7R^8$ ,  $-NR^6SO_2R^9$ ,  $-COOR^6$ ,  
 $-CONR^6R^7$ ,  $-COR^6$ ,  $-SO_2NR^6R^7$ ,  $-S(O)_{0-2}R^9$ ,  $-O(CH_2)_{1-10}-COOR^6$ ,  
 $-O(CH_2)_{1-10}CONR^6R^7$ ,  $-CF_3$ ,  $-CN$ ,  $-NO_2$  and halogen.

10 48. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 47 and a pharmaceutically acceptable carrier.

15 49. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 48.

20 50. A therapeutic combination comprising:

- (a) a first amount of at least one of nicotinic acid or derivatives thereof; and
- (b) a second amount of at least one sterol absorption inhibitor represented by Formula (V):



(V)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (V) or of the isomers thereof, or prodrugs of the compounds of Formula (V) or of the isomers, salts or solvates thereof, wherein, in Formula (V) above:

Ar<sup>1</sup> is aryl, R<sup>10</sup>-substituted aryl or heteroaryl;

5 Ar<sup>2</sup> is aryl or R<sup>4</sup>-substituted aryl;

Ar<sup>3</sup> is aryl or R<sup>5</sup>-substituted aryl;

X and Y are independently selected from the group consisting of -CH<sub>2</sub>-,  
-CH(lower alkyl)- and -C(dilower alkyl)-;

10 R is -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup> or -O(CO)NR<sup>6</sup>R<sup>7</sup>; R<sup>1</sup> is hydrogen, lower alkyl  
or aryl; or R and R<sup>1</sup> together are =O;

q is 0 or 1;

r is 0, 1 or 2;

m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and  
q is 1, 2, 3, 4 or 5;

15 R<sup>4</sup> is 1-5 substituents independently selected from the group consisting of  
lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>,  
-NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>,  
-CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>,  
-O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

20 R<sup>5</sup> is 1-5 substituents independently selected from the group consisting of -  
OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>R<sup>7</sup>,  
-NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>, -CONR<sup>6</sup>R<sup>7</sup>, -  
COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -  
NO<sub>2</sub>, halogen, -(lower alkylene)COOR<sup>6</sup> and -CH=CH-COOR<sup>6</sup>;

25 R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of  
hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;

R<sup>9</sup> is lower alkyl, aryl or aryl-substituted lower alkyl; and

30 R<sup>10</sup> is 1-5 substituents independently selected from the group consisting of  
lower alkyl, -OR<sup>6</sup>, -O(CO)R<sup>6</sup>, -O(CO)OR<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>6</sup>, -O(CO)NR<sup>6</sup>R<sup>7</sup>,  
-NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(CO)R<sup>7</sup>, -NR<sup>6</sup>(CO)OR<sup>9</sup>, -NR<sup>6</sup>(CO)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>SO<sub>2</sub>R<sup>9</sup>, -COOR<sup>6</sup>,


-CONR<sup>6</sup>R<sup>7</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>6</sup>,  
-O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>6</sup>R<sup>7</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen,

wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes,  
5 obesity or lowering a concentration of a sterol in plasma of a mammal.

51. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of  
10 the therapeutic combination of claim 50.

52. A composition comprising:

- (a) at least one of nicotinic acid or derivatives thereof; and  
15 (b) at least one sterol absorption inhibitor represented by Formula (VI):



(VI)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
20 of Formula (VI) or of the isomers thereof, or prodrugs of the compounds of Formula (VI) or of the isomers, salts or solvates thereof, wherein:

R<sub>1</sub> is

$\overset{|}{-CH-}$ ,  $\overset{|}{-C}(\text{lower alkyl})-$ ,  $\overset{|}{-CF-}$ ,  $\overset{|}{-C(OH)-}$ ,  $\overset{|}{-C(C_6H_5)-}$ ,  $\overset{|}{-C(C_6H_4-R_{15})-}$ ,  
 $\overset{|}{-N-}$  or  $\overset{|}{-N^+O^-}$ ;

25 R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of:  
-CH<sub>2</sub>-, -CH(lower alkyl)-, -C(di-lower alkyl)-, -CH=CH- and -C(lower alkyl)=CH-; or  
R<sub>1</sub> together with an adjacent R<sub>2</sub>, or R<sub>1</sub> together with an adjacent R<sub>3</sub>, form a



-CH=CH- or a -CH=C(lower alkyl)- group;

u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sub>2</sub> is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R<sub>3</sub> is -CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, the R<sub>2</sub>'s can be the same or different; and provided that when u is 2 or 3, the R<sub>3</sub>'s can be the same or different;

R<sub>4</sub> is selected from B-(CH<sub>2</sub>)<sub>m</sub>C(O)-, wherein m is 0, 1, 2, 3, 4 or 5;

B-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 0, 1, 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein Z is -O-, -C(O)-, phenylene, -N(R<sub>8</sub>)- or -S(O)<sub>0-2</sub>-, e is 0, 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4, 5 or 6;

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-;

B-(C<sub>4</sub>-C<sub>6</sub> alkadienylene)-;

B-(CH<sub>2</sub>)<sub>t</sub>-Z-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1, 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0, 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6;


B-(CH<sub>2</sub>)<sub>t</sub>-V-(C<sub>2</sub>-C<sub>6</sub> alkenylene)- or

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-V-(CH<sub>2</sub>)<sub>t</sub>-, wherein V and t are as defined above, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6; B-(CH<sub>2</sub>)<sub>a</sub>-Z-(CH<sub>2</sub>)<sub>b</sub>-V-(CH<sub>2</sub>)<sub>d</sub>-, wherein Z and V are as defined above and a, b and d are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3, 4, 5 or 6; or


T-(CH<sub>2</sub>)<sub>s</sub>-, wherein T is cycloalkyl of 3-6 carbon atoms and s is 0, 1, 2, 3, 4, 5 or 6; or

R<sub>1</sub> and R<sub>4</sub> together form the group  $\text{B}-\text{CH}=\overset{\text{I}}{\text{C}}-$  ;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl, thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or



- W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy, alkoxycarbonylalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl
- 5 lower alkanedioyl, allyloxy, -CF<sub>3</sub>, -OCF<sub>3</sub>, benzyl, R<sub>7</sub>-benzyl, benzyloxy, R<sub>7</sub>-benzyloxy, phenoxy, R<sub>7</sub>-phenoxy, dioxolanyl, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylenyloxy-, OH, halogeno, -CN, -N<sub>3</sub>, -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, R<sub>11</sub>O<sub>2</sub>SNH-, (R<sub>11</sub>O<sub>2</sub>S)<sub>2</sub>N-, -S(O)<sub>2</sub>NH<sub>2</sub>, -S(O)<sub>0-2</sub>R<sub>8</sub>, tert-butyl dimethyl-silyloxymethyl, -C(O)R<sub>12</sub>,
- 10 -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -CH=CHC(O)R<sub>12</sub>, -lower alkylene-C(O)R<sub>12</sub>, R<sub>10</sub>C(O)(lower alkylenyloxy)-, N(R<sub>8</sub>)(R<sub>9</sub>)C(O)(lower alkylenyloxy)- and



for substitution on ring carbon atoms,

- and the substituents on the substituted heteroaryl ring nitrogen atoms, when present, are selected from the group consisting of lower alkyl, lower alkoxy, C(O)OR<sub>10</sub>, -C(O)R<sub>10</sub>, OH, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-,
- 15 N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylenyloxy-, -S(O)<sub>2</sub>NH<sub>2</sub> and 2-(trimethylsilyl)-ethoxymethyl;


R<sub>7</sub> is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -COOH, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), OH, and halogeno;

R<sub>8</sub> and R<sub>9</sub> are independently selected from H or lower alkyl;

- 20 R<sub>10</sub> is selected from lower alkyl, phenyl, R<sub>7</sub>-phenyl, benzyl or R<sub>7</sub>-benzyl;

R<sub>11</sub> is selected from OH, lower alkyl, phenyl, benzyl, R<sub>7</sub>-phenyl or R<sub>7</sub>-benzyl;

R<sub>12</sub> is selected from H, OH, alkoxy, phenoxy, benzyloxy,



, -N(R<sub>8</sub>)(R<sub>9</sub>), lower alkyl, phenyl or R<sub>7</sub>-phenyl;

R<sub>13</sub> is selected from -O-, -CH<sub>2</sub>-, -NH-, -N(lower alkyl)- or -NC(O)R<sub>19</sub>;

- R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of H
- 25 and the groups defined for W; or R<sub>15</sub> is hydrogen and R<sub>16</sub> and R<sub>17</sub>, together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;

R<sub>19</sub> is H, lower alkyl, phenyl or phenyl lower alkyl; and


R<sub>20</sub> and R<sub>21</sub> are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzofused  
 5 heteroaryl, W-substituted benzofused heteroaryl and cyclopropyl, wherein heteroaryl is as defined above.

53. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma  
 10 of a mammal, comprising a therapeutically effective amount of the composition of claim 52 and a pharmaceutically acceptable carrier.

54. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the  
 15 step of administering to a mammal in need of such treatment an effective amount of the composition of claim 53.

55. A therapeutic combination comprising:


- (a) a first amount of at least one of nicotinic acid or derivatives thereof; and  
 20  
 (b) a second amount of at least one sterol absorption inhibitor represented by Formula (VI):



(VI)

25 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VI) or of the isomers thereof, or prodrugs of the compounds of Formula (VI) or of the isomers, salts or solvates thereof, wherein:

R<sub>1</sub> is



R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of:

- 5    -CH<sub>2</sub>-, -CH(lower alkyl)-, -C(di-lower alkyl)-, -CH=CH- and -C(lower alkyl)=CH-; or  
 R<sub>1</sub> together with an adjacent R<sub>2</sub>, or R<sub>1</sub> together with an adjacent R<sub>3</sub>, form a  
 -CH=CH- or a -CH=C(lower alkyl)- group;

u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that  
 when R<sub>2</sub> is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R<sub>3</sub> is -

- 10    CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, the R<sub>2</sub>'s can be  
 the same or different; and provided that when u is 2 or 3, the R<sub>3</sub>'s can be the same or  
 different;

R<sub>4</sub> is selected from B-(CH<sub>2</sub>)<sub>m</sub>C(O)-, wherein m is 0, 1, 2, 3, 4 or 5;

B-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 0, 1, 2, 3, 4, 5 or 6;

- 15    B-(CH<sub>2</sub>)<sub>e</sub>-Z-(CH<sub>2</sub>)<sub>r</sub>-, wherein Z is -O-, -C(O)-, phenylene, -N(R<sub>8</sub>)- or -S(O)<sub>0-2</sub>-, e is 0,  
 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4,  
 5 or 6;

B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-;

B-(C<sub>4</sub>-C<sub>6</sub> alkadienylene)-;

- 20    B-(CH<sub>2</sub>)<sub>t</sub>-Z-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1,  
 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene  
 chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0,  
 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6;

- 25    B-(CH<sub>2</sub>)<sub>t</sub>-V-(C<sub>2</sub>-C<sub>6</sub> alkenylene)- or


B-(C<sub>2</sub>-C<sub>6</sub> alkenylene)-V-(CH<sub>2</sub>)<sub>t</sub>-, wherein V and t are as defined above, provided that  
 the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6;

B-(CH<sub>2</sub>)<sub>a</sub>-Z-(CH<sub>2</sub>)<sub>b</sub>-V-(CH<sub>2</sub>)<sub>d</sub>-, wherein Z and V are as defined above and a, b and d  
 are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3,  
 4, 5 or 6; or


- 30    T-(CH<sub>2</sub>)<sub>s</sub>-, wherein T is cycloalkyl of 3-6 carbon atoms and s is 0, 1, 2, 3, 4, 5 or 6; or

R<sub>1</sub> and R<sub>4</sub> together form the group  $B-CH=\overset{|}{C}-$  ;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl,  
 5 thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or



W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy,  
 10 alkoxycarbonylalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl lower alkanedioyl, allyloxy, -CF<sub>3</sub>, -OCF<sub>3</sub>, benzyl, R<sub>7</sub>-benzyl, benzyloxy, R<sub>7</sub>-benzyloxy, phenoxy, R<sub>7</sub>-phenoxy, dioxolanyl, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, OH, halogeno, -CN, -N<sub>3</sub>, -NHC(O)OR<sub>10</sub>, -NHC(O)R<sub>10</sub>, R<sub>11</sub>O<sub>2</sub>SNH-, (R<sub>11</sub>O<sub>2</sub>S)<sub>2</sub>N-, -S(O)<sub>2</sub>NH<sub>2</sub>, -  
 15 S(O)<sub>0-2</sub>R<sub>8</sub>, tert-butyldimethyl-silyloxymethyl, -C(O)R<sub>12</sub>, -COOR<sub>19</sub>, -CON(R<sub>8</sub>)(R<sub>9</sub>), -CH=CHC(O)R<sub>12</sub>, -lower alkylene-C(O)R<sub>12</sub>, R<sub>10</sub>C(O)(lower alkyleneoxy)-, N(R<sub>8</sub>)(R<sub>9</sub>)C(O)(lower alkyleneoxy)- and



for substitution on ring carbon atoms,

and the substituents on the substituted heteroaryl ring nitrogen atoms, when present,  
 20 are selected from the group consisting of lower alkyl, lower alkoxy, -C(O)OR<sub>10</sub>, -C(O)R<sub>10</sub>, OH, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkylene-, N(R<sub>8</sub>)(R<sub>9</sub>)-lower alkyleneoxy-, -S(O)<sub>2</sub>NH<sub>2</sub> and 2-(trimethylsilyl)-ethoxymethyl;


R<sub>7</sub> is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -COOH, NO<sub>2</sub>, -N(R<sub>8</sub>)(R<sub>9</sub>), OH, and halogeno;

25 R<sub>8</sub> and R<sub>9</sub> are independently selected from H or lower alkyl;

R<sub>10</sub> is selected from lower alkyl, phenyl, R<sub>7</sub>-phenyl, benzyl or R<sub>7</sub>-benzyl;

R<sub>11</sub> is selected from OH, lower alkyl, phenyl, benzyl, R<sub>7</sub>-phenyl or R<sub>7</sub>-benzyl;

R<sub>12</sub> is selected from H, OH, alkoxy, phenoxy, benzyloxy,



, -N(R<sub>8</sub>)(R<sub>9</sub>), lower alkyl, phenyl or R<sub>7</sub>-phenyl;

R<sub>13</sub> is selected from -O-, -CH<sub>2</sub>-, -NH-, -N(lower alkyl)- or -NC(O)R<sub>19</sub>;

R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of H and the groups defined for W; or R<sub>15</sub> is hydrogen and R<sub>16</sub> and R<sub>17</sub>, together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;

R<sub>19</sub> is H, lower alkyl, phenyl or phenyl lower alkyl; and

R<sub>20</sub> and R<sub>21</sub> are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzofused heteroaryl, W-substituted benzofused heteroaryl and cyclopropyl, wherein heteroaryl is as defined above,


wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

56. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the therapeutic combination of claim 55.

57. A composition comprising:


(a) at least one of nicotinic acid or derivatives thereof; and

(b) at least one sterol absorption inhibitor represented by Formula (VIIA) or (VIIB):



(VIIA)

or




(VIIB)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds  
 5 of Formulae (VIIA) or (VIIB) or of the isomers thereof, or prodrugs of the compounds  
 of Formulae (VIIA) or (VIIB) or of the isomers, salts or solvates thereof, wherein in  
 Formulae (VIIA) and (VIIB):


A is -CH=CH-, -C≡C- or -(CH<sub>2</sub>)<sub>p</sub>- wherein p is 0, 1 or 2;

B is



10

B' is




D is -(CH<sub>2</sub>)<sub>m</sub>C(O)- or -(CH<sub>2</sub>)<sub>q</sub>- wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;

E is C<sub>10</sub> to C<sub>20</sub> alkyl or -C(O)-(C<sub>9</sub> to C<sub>19</sub>)-alkyl, wherein the alkyl is straight or  
 15 branched, saturated or containing one or more double bonds;

R is hydrogen, C<sub>1</sub>-C<sub>15</sub> alkyl, straight or branched, saturated or containing one  
 or more double bonds, or B-(CH<sub>2</sub>)<sub>r</sub>-, wherein r is 0, 1, 2, or 3;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>1</sub>', R<sub>2</sub>', and R<sub>3</sub>' are independently selected from the group  
 consisting of hydrogen, lower alkyl, lower alkoxy, carboxy, NO<sub>2</sub>, NH<sub>2</sub>, OH, halogeno,  
 20 lower alkylamino, dilower alkylamino, -NHC(O)OR<sub>5</sub>, R<sub>6</sub>O<sub>2</sub>SNH- and -S(O)<sub>2</sub>NH<sub>2</sub>;

R<sub>4</sub> is



wherein n is 0, 1, 2 or 3;

$R_5$  is lower alkyl; and


$R_6$  is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents  
 5 are 1-3 groups independently selected from the group consisting of lower alkyl, lower  
 alkoxy, carboxy,  $NO_2$ ,  $NH_2$ , OH, halogeno, lower alkylamino and dilower alkylamino.

58. A pharmaceutical composition for the treatment or prevention of a  
 vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma  
 10 of a mammal, comprising a therapeutically effective amount of the composition of  
 claim 57 and a pharmaceutically acceptable carrier.

59. A method of treating or preventing a vascular condition, diabetes,  
 obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the  
 15 step of administering to a mammal in need of such treatment an effective amount of  
 the composition of claim 58.

60. A therapeutic combination comprising:


- (a) a first amount of at least one of nicotinic acid or derivatives thereof; and  
 20  
 (b) a second amount of at least one sterol absorption inhibitor represented  
 by Formula (VIIA) or (VIIB):



(VIIA)

25 or






(VII B)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formulae (VIIA) or (VII B) or of the isomers thereof, or prodrugs of the compounds of Formulae (VIIA) or (VII B) or of the isomers, salts or solvates thereof, wherein in


5 of Formulae (VIIA) and (VII B):

A is -CH=CH-, -C≡C- or -(CH<sub>2</sub>)<sub>p</sub>- wherein p is 0, 1 or 2;

B is



10 B' is




D is -(CH<sub>2</sub>)<sub>m</sub>C(O)- or -(CH<sub>2</sub>)<sub>q</sub>- wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;

E is C<sub>10</sub> to C<sub>20</sub> alkyl or -C(O)-(C<sub>9</sub> to C<sub>19</sub>)-alkyl, wherein the alkyl is straight or branched, saturated or containing one or more double bonds;

15 R is hydrogen, C<sub>1</sub>-C<sub>15</sub> alkyl, straight or branched, saturated or containing one or more double bonds, or B-(CH<sub>2</sub>)<sub>r</sub>-, wherein r is 0, 1, 2, or 3;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>1</sub>', R<sub>2</sub>', and R<sub>3</sub>' are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, carboxy, NO<sub>2</sub>, NH<sub>2</sub>, OH, halogeno, lower alkylamino, dilower alkylamino, -NHC(O)OR<sub>5</sub>, R<sub>6</sub>O<sub>2</sub>SNH- and -S(O)<sub>2</sub>NH<sub>2</sub>;

20 R<sub>4</sub> is



wherein n is 0, 1, 2 or 3;

R<sub>5</sub> is lower alkyl; and


R<sub>6</sub> is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents are 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, carboxy, NO<sub>2</sub>, NH<sub>2</sub>, OH, halogeno, lower alkylamino and dilower alkylamino, wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

61. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the therapeutic combination of claim 60.

62. A composition comprising:

(a) at least one of nicotinic acid or derivatives thereof; and

(b) at least one sterol absorption inhibitor represented by Formula (VIII):





(VIII)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIII) or of the isomers thereof, or prodrugs of the compounds of Formula (VIII) or of the isomers, salts or solvates thereof, wherein, in Formula (VIII) above,

R<sup>26</sup> is H or OG<sup>1</sup>;

G and G<sup>1</sup> are independently selected from the group consisting of





provided that when R<sup>26</sup> is H or

OH, G is not H;

R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy or -W-R<sup>30</sup>;

W is independently selected from the group consisting of  
 5 -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and  
 -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group  
 10 consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and  
 -C(O)aryl;

R<sup>30</sup> is selected from the group consisting of R<sup>32</sup>-substituted T,  
 R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl,  
 R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and  
 15 R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl,  
 oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl,  
 imidazolyl and pyridyl;


20 R<sup>32</sup> is independently selected from 1-3 substituents independently selected  
 from the group consisting of halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy,  
 -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl,  
 (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and  
 25 pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is

attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

5 Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone,

forms the spiro group ; and

R<sup>1</sup> is selected from the group consisting of


10 -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

-(CH<sub>2</sub>)<sub>e</sub>-E-(CH<sub>2</sub>)<sub>r</sub>-, wherein E is -O-, -C(O)-, phenylene, -NR<sup>22</sup>- or -S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C<sub>2</sub>-C<sub>6</sub>)alkenylene-; and

15 -(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R<sup>12</sup> is



20 R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub> alkyl))-, -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;


provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1;

provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1;

25 provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and

provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

and when Q is a bond, R<sup>1</sup> also can be:



M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub>)alkyl- and -C(di-(C<sub>1</sub>-C<sub>6</sub>)alkyl);

- 5        R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>,  
 10        -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>15</sup> and R<sup>17</sup> are independently selected from the group consisting of -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup> and -O(CO)NR<sup>19</sup>R<sup>20</sup>;

- 15        R<sup>16</sup> and R<sup>18</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>15</sup> and R<sup>16</sup> together are =O, or R<sup>17</sup> and R<sup>18</sup> together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

- 20        s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

- 25        j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R<sup>1</sup> is 
$$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$$
, Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and


R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

63. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 62 and a pharmaceutically acceptable carrier.

64. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 63.

65. A therapeutic combination comprising:

- (a) a first amount of at least one of nicotinic acid or derivatives thereof; and
- (b) a second amount of at least one sterol absorption inhibitor represented by Formula (VIII):





(VIII)

or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (VIII) or of the isomers thereof, or prodrugs of the compounds of Formula (VIII) or of the isomers, salts or solvates thereof, wherein, in Formula (VIII) above,

$R^{26}$  is H or  $OG^1$ ;

G and  $G^1$  are independently selected from the group consisting of



provided that when  $R^{26}$  is H or

OH, G is not H;

R,  $R^a$  and  $R^b$  are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy or -W- $R^{30}$ ;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N( $R^{31}$ )-, -NH-C(O)-N( $R^{31}$ )- and -O-C(S)-N( $R^{31}$ )-;

$R^2$  and  $R^6$  are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

$R^3$ ,  $R^4$ ,  $R^5$ ,  $R^7$ ,  $R^{3a}$  and  $R^{4a}$  are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

$R^{30}$  is selected from the group consisting of  $R^{32}$ -substituted T,  $R^{32}$ -substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl,  $R^{32}$ -substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl,

R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and  
 R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;


T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl,  
 5 oxazolyl, isoxazolyl, thiazolyl, iosthiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl,  
 imidazolyl and pyridyl;

R<sup>32</sup> is independently selected from 1-3 substituents independently selected  
 from the group consisting of halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy,  
 -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl,  
 10 (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and  
 pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is  
 attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolyl or  
 morpholyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-  
 15 methylpiperazinyl, indolyl or morpholyl group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone,

forms the spiro group ; and

20 R<sup>1</sup> is selected from the group consisting of

-(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, provided that when Q forms a spiro ring, q  
 can also be zero or 1;


-(CH<sub>2</sub>)<sub>e</sub>-E-(CH<sub>2</sub>)<sub>r</sub>-, wherein E is -O-, -C(O)-, phenylene, -NR<sup>22</sup>- or -  
 S(O)<sub>0-2</sub>-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

25 -(C<sub>2</sub>-C<sub>6</sub>)alkenylene-; and

-(CH<sub>2</sub>)<sub>f</sub>-V-(CH<sub>2</sub>)<sub>g</sub>-, wherein V is C<sub>3</sub>-C<sub>6</sub> cycloalkylene, f is 1-5 and g is 0-  
 5, provided that the sum of f and g is 1-6;

R<sup>12</sup> is





R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub>) alkyl)-, -CH=CH- and

-C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;


provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1;

provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1;

provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and

provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

and when Q is a bond, R<sup>1</sup> also can be:



M is -O-, -S-, -S(O)- or -S(O)<sub>2</sub>-;

X, Y and Z are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub>)alkyl- and -C(di-(C<sub>1</sub>-C<sub>6</sub>)alkyl);

R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>, -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

R<sup>15</sup> and R<sup>17</sup> are independently selected from the group consisting of -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup> and -O(CO)NR<sup>19</sup>R<sup>20</sup>;

R<sup>16</sup> and R<sup>18</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl and aryl; or R<sup>15</sup> and R<sup>16</sup> together are =O, or R<sup>17</sup> and R<sup>18</sup> together are =O;

d is 1, 2 or 3;

5 h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;

10 v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R<sup>1</sup> is 
$$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j-(\text{C})_v-\text{Y}_k-\text{S}(\text{O})_{0-2}- \\ | \\ \text{R}^{16} \end{array}$$
, Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

15 R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

20 R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy,


25 wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

66. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of  
30 the therapeutic combination of claim 65.

67. A composition comprising:

(a) at least one of nicotinic acid or derivatives thereof; and

5 (b) at least one sterol absorption inhibitor represented by Formula (IX):




10 or isomers thereof, or pharmaceutically acceptable salts or solvates of the compounds of Formula (IX) or of the isomers thereof, or prodrugs of the compounds of Formula (IX) or of the isomers, salts or solvates thereof, wherein in Formula (IX):

R<sup>26</sup> is selected from the group consisting of:

- 15 a) OH;  
b) OCH<sub>3</sub>;  
c) fluorine and  
d) chlorine.

R<sup>1</sup> is selected from the group consisting of



R, R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of H, -OH, halogeno, -NH<sub>2</sub>, azido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)-alkoxy and -W-R<sup>30</sup>;

20 W is independently selected from the group consisting of

-NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2</sup> and R<sup>6</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>3a</sup> and R<sup>4a</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl and -C(O)aryl;

R<sup>30</sup> is independently selected from the group consisting of R<sup>32</sup>-substituted T, R<sup>32</sup>-substituted-T-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, R<sup>32</sup>-substituted-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and R<sup>32</sup>-substituted-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>31</sup> is independently selected from the group consisting of H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;


T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R<sup>32</sup> is independently selected from 1-3 substituents independently selected from the group consisting of H, halogeno, (C<sub>1</sub>-C<sub>4</sub>)alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, methylenedioxy, oxo, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-N((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>2</sub>, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkoxy and pyrrolidinylcarbonyl; or R<sup>32</sup> is a covalent bond and R<sup>31</sup>, the nitrogen to which it is attached and R<sup>32</sup> form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar<sup>1</sup> is aryl or R<sup>10</sup>-substituted aryl;

Ar<sup>2</sup> is aryl or R<sup>11</sup>-substituted aryl;

Q is -(CH<sub>2</sub>)<sub>q</sub>-, wherein q is 2-6, or, with the 3-position ring carbon of the azetidinone,



R<sup>12</sup> is

$\overset{|}{-}\text{CH}-$ ,  $\overset{|}{-}\text{C}(\text{C}_1\text{-C}_6 \text{ alkyl})-$ ,  $\overset{|}{-}\text{CF}-$ ,  $\overset{|}{-}\text{C}(\text{OH})-$ ,  $\overset{|}{-}\text{C}(\text{C}_6\text{H}_4\text{-R}^{23})-$ ,  $\overset{|}{-}\text{N}-$ , or  $\overset{|}{-}\text{NO}^+$ ;

R<sup>13</sup> and R<sup>14</sup> are independently selected from the group consisting of -CH<sub>2</sub>-, -CH(C<sub>1</sub>-C<sub>6</sub> alkyl)-, -C(di-(C<sub>1</sub>-C<sub>6</sub> alkyl)), -CH=CH- and -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-; or R<sup>12</sup> together with an adjacent R<sup>13</sup>, or R<sup>12</sup> together with an adjacent R<sup>14</sup>, form a -CH=CH- or a -CH=C(C<sub>1</sub>-C<sub>6</sub> alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R<sup>13</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, a is 1; provided that when R<sup>14</sup> is -CH=CH- or -C(C<sub>1</sub>-C<sub>6</sub> alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R<sup>13</sup>'s can be the same or different; and provided that when b is 2 or 3, the R<sup>14</sup>'s can be the same or different;

R<sup>10</sup> and R<sup>11</sup> are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, -OR<sup>19</sup>, -O(CO)R<sup>19</sup>, -O(CO)OR<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-5</sub>OR<sup>19</sup>, -O(CO)NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>R<sup>20</sup>, -NR<sup>19</sup>(CO)R<sup>20</sup>, -NR<sup>19</sup>(CO)OR<sup>21</sup>, -NR<sup>19</sup>(CO)NR<sup>20</sup>R<sup>25</sup>, -NR<sup>19</sup>SO<sub>2</sub>R<sup>21</sup>, -COOR<sup>19</sup>, -CONR<sup>19</sup>R<sup>20</sup>, -COR<sup>19</sup>, -SO<sub>2</sub>NR<sup>19</sup>R<sup>20</sup>, S(O)<sub>0-2</sub>R<sup>21</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>-COOR<sup>19</sup>, -O(CH<sub>2</sub>)<sub>1-10</sub>CONR<sup>19</sup>R<sup>20</sup>, -(C<sub>1</sub>-C<sub>6</sub> alkylene)-COOR<sup>19</sup>, -CH=CH-COOR<sup>19</sup>, -CF<sub>3</sub>, -CN, -NO<sub>2</sub> and halogen;

Ar<sup>1</sup> can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R<sup>19</sup> and R<sup>20</sup> are independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and aryl-substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or R<sup>24</sup>-substituted aryl;

R<sup>22</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl (C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)R<sup>19</sup> or -COOR<sup>19</sup>;

R<sup>23</sup> and R<sup>24</sup> are independently 1-3 groups independently selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -COOH, NO<sub>2</sub>, -NR<sup>19</sup>R<sup>20</sup>, -OH and halogeno; and

R<sup>25</sup> is H, -OH or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

68. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 67 and a pharmaceutically acceptable carrier.

5

69. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 68.

10

70. A composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one substituted azetidinone compound or a pharmaceutically acceptable salt or solvate thereof, or prodrug of the at least one substituted azetidinone compound or of the isomers, salts or solvates thereof.

15

71. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 70 and a pharmaceutically acceptable carrier.

20

72. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 70.

25

73. A therapeutic combination comprising:

(a) a first amount of at least one of nicotinic acid or derivatives thereof; and

(b) a second amount of at least one substituted azetidinone compound or a pharmaceutically acceptable salt or solvate thereof, or prodrug of the at least one substituted azetidinone compound or of the isomers, salts or solvates thereof,

30

wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

5           74. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the therapeutic combination of claim 73 and a pharmaceutically acceptable carrier.

10           75. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 73.

15           76. A composition comprising: (a) at least one of nicotinic acid or derivatives thereof; and (b) at least one substituted  $\beta$ -lactam compound or a pharmaceutically acceptable salt or solvate thereof, or prodrug of the at least one substituted  $\beta$ -lactam compound or of the isomers, salts or solvates thereof.

20           77. A pharmaceutical composition for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the composition of claim 76 and a pharmaceutically acceptable carrier.

25           78. A method of treating or preventing a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 77.

30           79. A therapeutic combination comprising:  
          (a) a first amount of at least one of nicotinic acid or derivatives thereof; and

- (c) a second amount of at least one substituted  $\beta$ -lactam compound or a pharmaceutically acceptable salt or solvate thereof, or prodrug of the at least one substituted  $\beta$ -lactam compound or of the isomers, salts or solvates thereof,
- 5 wherein the first amount and the second amount together comprise a therapeutically effective amount for the treatment or prevention of a vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal.

80. A pharmaceutical composition for the treatment or prevention of a  
10 vascular condition, diabetes, obesity or lowering a concentration of a sterol in plasma of a mammal, comprising a therapeutically effective amount of the therapeutic combination of claim 79 and a pharmaceutically acceptable carrier.

81. A method of treating or preventing a vascular condition, diabetes,  
15 obesity or lowering a concentration of a sterol in plasma of a mammal, comprising the step of administering to a mammal in need of such treatment an effective amount of the composition of claim 80.